

Biodegradation of Diesel by Aeromonashydrophila

Lakshmi Priya .J

K.S.Rangasamy College of Arts and Science, Tiruchengode/ Periyar University/India

ABSTRACT: *Hydrocarbon degrading microorganisms play a major role in the environment.. The purpose of the present study was to enumerate of Aeromonas sp from oil contaminated soil and to study degrading capacity, emulsification activity and production of biosurfactant. The hydrocarbon substrate specificity test shown that diesel is also one of the best substrate for growth and emulsification of biosurfactant by Aeromonas. Among 6 strains of Aeromonas, first strain(A1) shows maximum degradation rate at end of 168 hrs upto 19.37% followed by other strains, about 75% of diesel was degraded by Aeromonas over a period of 7 days. Emulsification upto 75% by A1 followed by A2(63.75%), A3(57.5%), Biosurfactant production by A1 strain 0.064g/l followed by other strains it represents a new type of biosurfactant with strong emulsifying ability.*

Keywords: *Aeromonas, Biosurfactant, Diesel, Emulsify, Soil*

I. INTRODUCTION

Biodegradation is the partial or complete conversion of the compound of interest to its elements. The role of organisms, both micro-and macro-organisms, in biodegradation is complex. It is a function of the organism's presence in an environment, their ecology, their metabolism (enzyme complement and efficiencies), growth rate and kinetics (of both growth and metabolism). It has been known for several decades that microorganisms possess both aerobic and anaerobic degradation. Microorganisms are actively involved in the degradation of several naturally occurring and toxic substances such as petroleum hydrocarbons, pesticides etc., Aerobic biodegradation is the breakdown of organic contaminants by microorganisms when oxygen is present. Aerobic bacteria use oxygen as an electron acceptor, and break down organic chemicals into smaller or organic compounds, often producing carbon dioxide and water as the final product. Aerobic biodegradation is also known as aerobic respiration. Aerobic biodegradation is an important component of the natural attenuation of contaminants at many hazardous waste sites. Anaerobic biodegradation is the breakdown of organic contaminants by microorganisms when oxygen is not present. Some anaerobic bacteria use nitrate, sulfate, iron, manganese, and carbon dioxide as their electron acceptors. And break down organic chemicals into smaller compounds, often producing carbon dioxide and methane as the final products. Anaerobic biodegradation is an important components of the natural attenuation of contaminants at many hazardous waste sites.

The most striking feature of a survey of the microorganisms involved in biodegradation processes is their large numbers, ubiquitous presence and varied capabilities. Rather than list the organisms involved (including the bacteria, fungi, actinomycetes, protozoa, etc.) a very brief treatment of the reasons for this versatility is given. Microorganisms as a group show a very wide tolerance range for environmental factors – very low to very high pH levels, 0°C to 80°C temperature, of the reasons for this versatility is given.

Biodegradation of organic compounds is the partial simplification or complete destruction of their molecular structure by physiological reactions catalyzed by microorganisms (Alexander, 1981; Atlas and Bartha, 1992; Young, 1997). Biodegradation is routinely measured by applying chemical and physiological assays to laboratory incubations of flasks containing pure cultures of microorganisms, mixed cultures, or environmental samples (e.g. soil, water, sediment, or industrial sludges). Oxygen is one of the most important requirements for microbial degradation of hydrocarbons. However, its availability is rarely a rate-limiting factor in the biodegradation of marine oil spills. Microorganisms employ oxygen-incorporating enzymes to initiate attack on hydrocarbons. Anaerobic degradation of certain hydrocarbons (i.e., degradation in the absence of oxygen) also occurs, but usually at negligible rates. Such degradation follows different chemical paths, and its ecological significance is generally considered minor. Studies of sediments impacted by the Amoco Cadiz spill found that, at best, anaerobic biodegradation is several orders of magnitude slower than aerobic biodegradation (Ward et al., 1980).

1.1 Hydrocarbons and its impact on the environment

Petroleum hydrocarbons existed long before humans developed the technological ability to retrieve it from the earth and use it as a source of energy. Natural seeps within the ocean floor have been releasing the

hydrocarbons for thousands of years creating ecosystems with adaptive microorganisms that utilize petroleum effectively. However, the ecological balance in environments that are not adjusted to assimilating large amount of spills or released from large quantity transportation and extraction practices.

Since the mid-1980's, hydrocarbon contamination has become a critical environmental issue in the world due its adverse environmental and health effects. Hence, increasing attention is being given to the study and development of techniques for cleaning up this contamination. To understand the potential environmental impact that can occur during the oil spill, the molecular components of petroleum hydrocarbons must be considered. The natural composition of petroleum diesel is very complex.

Petroleum hydrocarbon continues to be used as the principle source of energy and hence an important global environmental pollutant. Apart from accidental contamination of ecosystem, the vast amounts of oil sludge generated in refineries from water oil separation system and accumulation of waste oily materials in crude oil storage tank bottoms pose great problems because of the expensive disposal methods. (Ferrari et al., 1996; Vasudevan and Rajaram, 2001). Despite decades of research, successful bioremediation of petroleum hydrocarbon contaminated soil remains challenge.

Concentration of inorganic nutrients often limits the biodegradation of petroleum hydrocarbons in marine environment. Atlas and Bartha (1972) found that microbial degradation and mineralization were not increased by nitrate or phosphate alone but were increased dramatically when nitrate and phosphate were added together.


1.2 Distribution of hydrocarbon degrading microorganisms

It has been known for several years that certain microorganisms are able to degrade petroleum hydrocarbons and use them as sole source of carbon and energy for growth. The ability to degrade and/or utilize hydrocarbons substrates is exhibited by wide variety of bacterial genera, 25 genera of hydrocarbon degrading bacteria (Floodgate, 1984) have been isolated from the marine environment. Similarly 22 genera of hydrocarbon degrading bacteria have been reported (Bossert and Bartha, 1984) based on the number of published.

Microbial remediation of toxic hydrocarbon contaminated sites is carried out by a diverse group of microorganisms. Study of this diversity at the genetic level is necessary to understand the phylogenetic perspective, the mechanism of degradation, and develop novel strategies of treatment. Analysis of microorganisms having high specificity for recalcitrant compound. Documentation of this microbial diversity from oily sludge/crude oil contaminated sites is essential because they create a major environmental concern and these microbes can be used for cleaning up the same ().

Microbial degradation of oil has been shown to occur by attack on aliphatic or light aromatic fractions of the oil, with high molecular weight aromatics, resins, and asphaltene considered to be recalcitrant or exhibiting only very low rates of biodegradation (Joseph et al., 1990) Broderick and Coony (1982) reported that 96% of hydrocarbon utilizing bacteria isolated from freshwater lakes were able to emulsify kerosene, and it has been observed that mixed cultures of marine and soil bacteria which effectively degrade crude oil also exhibit strong emulsifying activity. Both aerobic and anaerobic biodegradation have been shown to reduce the concentration of several components of petroleum hydrocarbons. Mixed cultures often involve significant degradative capabilities because the single strains can complement to one another due to their physiological properties. Therefore, some members of the culture might be able to provide important degradative enzymes whereas other supply surfactants or growth factors.

Degradation pathway for hydrocarbons



During oil biodegradation, oil fluid properties change because different classes of compounds in petroleum have different susceptibilities to biodegradation. The early stages of oil biodegradation are characterized by loss of n-paraffin followed by loss of acyclic isoprenoids. Compared with those compounds groups, other compound classes are more resistant to biodegradation. However, even those more resistant compound classes are eventually destroyed as biodegradation proceeds. Hydrocarbons are natural products as well as pollutants; it is not surprising that hydrocarbon oxidizing bacteria, fungi and algae are distributed widely in nature. A typical soil, beach sand or ocean sediment contains 10^4 to 10^6 hydrocarbon degrading microorganisms per gram of soil (Rosenberg, 1991).

1.3 Characteristics of diesel

One can obtain diesel from petroleum, which is sometimes called petrodiesel when there is a need to distinguish it from diesel obtained from other sources. As a hydrocarbon mixture, it is obtained in the fractional distillation of crude oil between 250^oC and 350^oC at atmospheric pressure. Diesel is generally simpler to refine than gasoline and often costs less (through price fluctuations often mean that the inverse is true).

Reducing the level of sulfur in diesel is better for the environment. It allows the use of catalytic diesel particulate filters to control diesel particulate emissions, as well as more advanced technologies, such as Nox absorbers (still under development), to reduce emissions of nitrogen oxides (Nox). However, lowering sulfur also reduces the lubricity of the fuel, meaning that additives must be put into the fuel to help lubricate engines. Diesel contains approximately 18% more energy per unit of volume than gasoline, which along with the greater efficiency of diesel engines contributes of fuel economy (distance traveled per volume of fuel consumed). In the maritime field various grades of diesel fuel are used.

1.4 Role of Aeromonas in the degradation of hydrocarbons

Aeromonas is a non motile, gram negative rod shaped bacteria, chemoorganotrophic facultative anaerobes demonstrating both respiration and fermentative metabolism. Although it is a pathogen to human as well as fish, 29.6% of total population of Aeromonas were reported in soil followed by Pseudomonas and Bacillus.

Like Pseudomonas and Bacillus sp. Aeromonas also play a vital role in hydrocarbon degradation and production of biosurfactant. The peak growth and biosurfactant production was on the 8th day (Ilori et al., 2005). Hydrocarbon degrading microorganisms play a major role in the environment. Crude oil degrading bacterial strains were isolated from refinery oil contaminated soil which includes the Aeromonas sp. the rate of degradation at the 8th day 53.55% (Vivekanandhan et al., 1999)

Several studies on production of biosurfactants by Aeromonas sp. were also reported (Desai et al., 1987; Lang et al., 1987; Rosenberg, 1986; Wilkinson et al., 1985).

1.5 Biosurfactants

1.5.1 Classification of biosurfactants

Biosurfactants can be classified in several broad groups: Glycolipids, lipopeptides, lipopolysaccharides, phospholipids, fattyacids, and neutral lipids. The classification of biosurfactants is based on their chemical nature. Low molecular weight substances (e.g., rhamnolipid phospholipids, peptides), polymeric materials (e.g., proteins, polysaccharides) or particulate compounds (e.g., extracellular vesicles or microbial cells) (Rosenberg, 1986)

1.5.2 Production of biosurfactants by microorganisms

Biosurfactants, which are natural emulsifiers of hydrocarbons, are produced by some bacteria, fungi and yeast. Biosurfactant is defined as a surface active molecule containing both hydrophobic and hydrophilic components which is produced by microorganisms.

Biosurfactants can improve the bioavailability of hydrocarbons to the microbial cells by increasing the area at the aqueous hydrocarbon interface. This increases the rate of hydrocarbon dissolution and thereby utilization by microorganism (Gerson, 1993). Surface active biosurfactants are employed for enhanced oil recovery (Hart, 1989) and as flocculating agents, as detergents and adhesives (Zajie and Saffens, 1984).

1.5.3 Application of bio-surfactants

Research in the area of biosurfactants has expanded quite a lot in recent years due to its potential use in different areas, such as the food industry; agricultural, pharmaceutical, oil industry, neurochemistry and the paper and pulp industry. The development of this line of research is of paramount importance, mainly in view of

the present concern with protection of the environment. Therefore, the most significant advantage of microbial surfactants over chemical surfactants is its ecological acceptance because of its biodegradability and nontoxic to natural environments. The emphasis to date has been on enhanced oil recovery, cleaning oil spills, oil emulsification and in breaking industrially derived oil-in-water and water-in-oil emulsions. Dispersion and solubilization of organic compounds having low water solubility is an important step in bioremediation. Biosurfactants offer potential for dealing with this problem by increasing the dispersion and solubilization of organic compounds having limited water solubility.

Biosurfactants have been used for gene transfection, as ligands for binding immunoglobulins, as adjuvants or antigens and also as inhibitor for fibrin clot formation and activators of fibrin clot lysis. Biosurfactants have the potential to be used as a preventive strategy to delay the on set of pathogenic biofilm growth on catheters and other medical insertional materials, thus lowering the large number of hospital infections without the use of synthetic drugs and chemicals. In spite of the immense potential of the biosurfactants, their use still remains limited, because of their comparatively high production cost, as well as scant information on their toxicity towards human systems. However, it is only a matter of time before the full potential of biosurfactants is fully exploited and used in medical science (Karnath et al., 1999).

II. IMPORTANCE OF THE PRESENT INVESTIGATION

Petroleum hydrocarbon continues to be used as the principle source of energy. Wide scale production, transport use and disposal and petroleum globally have made it a major containment in both prevalence and quantity in the environment. Biosurfactant are a group of surface-active molecules produced mainly by hydrocarbon degrading microorganism it can degrade or transform the components of petroleum products. They are non-toxic, non-hazardous, Biodegradable and environmentally friendly components. Hence, reclamation of petroleum hydrocarbon polluted sites can be carried out by bioremediation, which is can enhance natural process of biodegradation using biosurfactant producing and oil degrading bacterial cultures. Bioremediation technologies generally aim at providing favourable conditions of certain, temperature and nutrients to enhance biological hydrocarbon break down.

Most work on biosurfactant production by microorganisms has been concentrated on the determination of the distinct polar moieties used for categorization into classes such as glycolipids and lipoproteins. While biosurfactant production by organisms such as *Pseudomonas* sp. *Achinetobacter* sp. *Bacillus* sp. *Rhodococcus* sp. and *Arthobacter* sp. have been well studied, only a few reports exist on the ability of *Aeromonas* sp. to the best of our knowledge, to produce biosurfactant when grown on hydrocarbon. Among the soil microorganisms which includes *Bacillus* and *Pseudomonas*, the presence of *Aeromonas* ranges about 29%. Apart from its pathogenicity in humans and fish, its also play a vital role in hydrocarbon degradations.

In the case of above mentioned discussion the present study is designed to carry out the following objectives.

1. To enumerate of *Aeromonas* sp. from oil contaminated soil.
2. To study the degradation capacity of *Aeromonas* sp.
3. To study the emulsification activity.
4. To estimate the biosurfactant production.

III. RESULTS AND DISCUSSION

Enumeration of bacterial population

The total heterotrophic bacterial population (THB) from oil contaminated soil samples was enumerated. After the enrichment process, 1ml of enriched culture was serially diluted and plated using pour plate technique. The THB population was ranged between 1.6×10^5 and 3.4×10^6 cfu/g whereas in uncontaminated soil, the THB population may vary between $10^7 \sim 10^9$ cfu/g. This variation is due to the presence of hydrocarbon in the contaminated soil. As per the studies of Rahman et al (2003) the size of the microbial population decrease is based on the chemical composition of contaminating oil and species of microorganisms present in the microbial community of particular ecosystem. Plate counts confirm the presence of a significant number of hydrocarbon-oxidizing organisms in soil. Morphologically different isolates were observed on the nutrient agar plates.

As it was aimed to work on the degradative potential of *Aeromonas* species, oxidase and catalase positive isolates were subjected to biochemical analysis. Gram-negative rod, motile, urease negative, it utilized lysine, citrate and produced gas from glucose. Maltose and lactose were not utilized. The organism reduced

nitrate, utilized ornithine, but not sorbitol. It was therefore putatively classified as the species of *Aeromonas*, taken for the present investigation.

In unpolluted ecosystem, hydrocarbon utilizers generally constitute less than 0.1% of the microbial community and in oil polluted ecosystems they can constitute upto 100% of the viable microorganisms. The microbial populations quantitatively reflect the degree or extent of exposure of that ecosystem to hydrocarbon contamination (Atlas 1981, Al- Gounaim et al., 1995).

The rapidly expanding literature on the oxidation and assimilation of hydrocarbon substrates by soil microorganisms attests to the widespread occurrence and ease of isolating these organisms from nature (McClay et. al., 2000 and Van Dyke, 1991). Population levels of hydrocarbon utilizers and their population within the microbial community appear to be a sensitive index of environmental exposure to hydrocarbons.

Microorganisms are known to attack specific compounds present in crude oil that is a complex mixture of saturates, aromatics and polar compounds (Bharathi and Vasudevan, 2001). An effective degradation of crude oil would require simultaneous action of several metabolically versatile microorganisms with favorable environmental conditions such as pH, temperature and availability of nutrients (Venkateswaran and Harayama, 1995). An oil spill in the environment leads to an adaptive process and if metabolically active hydrocarbon utilizing microorganisms could respond would be reduced considerably. The necessity for seeding with complementary population may not be capable of degrading a wide range of potential substrates in a complex mixture such as crude oil (Chhatre et al., 1996).

Growth substrate range determination

The results of the hydrocarbon substrate specificity test revealed that the organism had good growth on diesel as substrate. When no difference was noticed in the turbidity of the test flask and that of the control, it was taken as no growth (-), when slight increase in turbidity was noticed, it was taken as poor growth (+). Significant increase in turbidity was taken as good growth (++).

In the present study among six strains of *Aeromonas*, four strains shows increase in turbidity in the presence of diesel as substrate was taken as good growth (++) and two strains shows slight turbidity was taken as poor growth (+). Thus the substrate specificity test conclude that diesel also act as best substrates for growth of hydrocarbon degraders (Table 1).

The addition of hydrocarbons to an ecosystem may result in a selective increase in microorganisms capable of utilizing the hydrocarbons and those that are capable of utilizing metabolites produced by the hydrocarbon-utilisers (Venkateswaran and Harayanma, 1995, Ferrari et al., 1996). The enhancement or reduction will depend upon the chemical composition of the contaminating hydrocarbons and on the species of microorganisms present within the microbial community of the particular ecosystem (Atlas, 1995).

The added oil enriched for the species that have inherent petroleum hydrocarbon assimilating potential (Bossert and Bartha, 1984), whereas the less adapted species among the total heterotrophic population are gradually eliminated, resulting in qualitative shifts in species composition (Amadi, 1990).

Degradation of diesel

Hydrocarbon degradation by microbial communities depends on the composition of the community and its adaptive response to the presence of petroleum hydrocarbon. The organism used in this study was isolated from oil contaminated soil samples from petrol station at Tiruchengode, the organism therefore might have had prior exposure to hydrocarbons like diesel, thus it shows good growth in diesel.

In the present study the bacterial strain *Aeromonas* developed from oil polluted sites grown well on diesel. The rate of degradation is proportional to the bacterial population. The degradation of diesel was observed at every 24hrs interval and there was a corresponding increase in the bacterial cell population. The degradation of diesel after 24hrs was 2.57% at the end of 168 hrs strain A1 degraded upto 19.37%, shows maximum degradation followed by the other strains (Table 2 and Figure 1).

The rate of microbial degradation of hydrocarbons in soils is affected by several physico-chemical and biological parameters including the number and species of microorganisms present, the conditions for microbial degradation activity (e.g. presence of nutrient, oxygen, pH and temperature) the quality, quantity and bioavailability of the contaminants; and the soil characteristics such as particle size distribution (Margesin and Schinner, 1997).

Isolation of degrading strains was performed with diesel oil, crude oil as sole carbon source, the result showed a reduction of 30% while this value decreased to 0.15% in uncontaminated samples, which includes the strains of *Aeromonas* sp and *Pseudomonas vasicularis* (Saagua et al., 2002).

Petrodiesel oil degraders belong to the genera *Micrococcus*, *Corynebacterium*, *Bacillus*, *Enterobacter*, *Pseudomonas*, *Alcaligenes*, *Flavobacterium*, *Moraxella*, *Aeromonas*, *Acinetobacter* and *Vibrio*. The flora reflects the normal heterotrophic bacteria present in soil and native genera seem to be crude oil utilizers. Several other workers also reported on the above genera as hydrocarbon degrading microorganisms (Atlas 1981, Leahy and Colwell 1990, Banat et al., 2001).

The rate of hydrocarbon degradation ranged from 0 $\mu\text{g g}^{-1}$ to 0.60 $\mu\text{g g}^{-1}$ for control oil, 0.05 $\mu\text{g g}^{-1}$ to 1.67 gasoline and from 0.12 $\mu\text{g g}^{-1}$ diesel oil to 1.31 $\mu\text{g g}^{-1}$ for hydrocarbon contaminated soils respectively (Obire, 2001). The biodegradability of seven different crude oils was found to be highly dependent on their composition and on incubation temperature. At 20°C lighter oils had greater abiotic losses and were more susceptible to biodegradation than heavier oils (Atlas, 1995).

Chhatre et al., (1996) reported about 60% of degradation of crude oil using semicontinuous crude oil fed reactor using a four members consortium. Several other workers (Venkateswaran and Harayama 1995, Lal and Khanna, 1996, Sugiura et al., 1997) showed that a bacterial consortium was able to degrade 28-51% of saturate and 0-18% of aromatics present in crude oil or up to 60% crude oil by mixed consortia. The percentage of biodegradation was significantly higher than that achieved by individual isolates.

By the addition of metabolically active hydrocarbon utilizing microorganisms, the lag period before the indigenous microbial population respond to the addition of a complex mixture such as diesel oil can be reduced considerably (Del'Arco and De Franca, 1999; Bharathi and Vasudevan, 2001). Several other workers (Chhatre et al., 1996; Sugiura et al., 1997; Vasudevan and Rajaram, 2001) have described the ability of mixed bacterial consortia to degrade 28-51% of saturates and 0-18% of aromatics present in petrodiesel or upto 60% petrodiesel.

Bioaugmentation and bioaugmentation on the degradation of total petroleum hydrocarbon (TPH) in soils contaminated by diesel oil, showed the greatest degradation upto 72.7% by the number of diesel oil degrading microorganisms includes *Aeromonas*, *Bacillus*, *Acinetobacter* (Fatima et al., 2003).

A survey of soils from the northwest area of Canada for the presence of oil-utilizing microorganisms indicated that not all soils have an indigenous population capable of utilizing oil degradation rates of the bacterial consortium such as *Flavobacterium* and *Cytophaga* (41%), *Pseudomonas* sp. (34%), *Xanthomonas* (9%), *Aeromonas* (17%) respectively (Jobson et al., 1972).

Emulsification

The formation of a water-in-oil mixture. An emulsified mixture of water in oil is commonly called mousse. The presence of mousse indicates that a spill has been on the water for some time.

The liquid aromatic hydrocarbons were particularly not good substrates for emulsification, however diesel was found to be good substrates for emulsification in the present study out of 6 strains of *Aeromonas* A1, A2, A3 shows 75%, 57.5%, 63.75%, respectively found to be good emulsifier (Table 4 and Figure 3).

Rosenberg (1994) suggested that the natural role of emulsions is to enhance the growth of bacteria on petroleum hydrocarbons. The ability of the extracellular emulsifying agent of *Arthrobacter* sp, *Aeromonas* sp, *Pseudomonas* sp. has been reported (Rosenberg et al., 1979). Emulsification is known to enhance hydrocarbon metabolism (Berg et al., 1990; Hommel, 1993). Stability of emulsion in the presence of salt has been reported as one of the properties of the biosurfactant produced by *Bacillus licheniformis* strain (McInerney et al., 1990). Broderick and Cooney (1982) reported that 96% of hydrocarbon utilizing bacteria isolated from lakes were able to emulsify kerosene which effectively activity.

Rosenberg et al., (1991) reported that the ability of the extracellular emulsifying agent of *Arthrobacter* sp, *Aeromonas* and *Bacillus* sp. to emulsify crude oil and fractions of crude oil, is, gas oil was a better substrate induced emulsification than kerosene. In fact, emulsions of gas oil were as stable as crude-oil emulsions. Diesel light petroleum has yielded emulsions and the emulsions derived from kerosene and gasoline were unstable. Pentane and hexane also were not emulsified effectively; however quantitative data for these two paraffins were not obtained because of extensive evaporation during incubation. A higher emulsifying activity has been reported that the biosurfactant produced by *Pseudomonas*. The emulsions were stable at temperatures ranging from 0°C to 100°C (Rosenberg, 1992).

Most microbial surfactants are substrate specific, solubilizing or emulsifying different hydrocarbons at different rates (Ilori and Amund, 2001). An emulsion is formed when one liquid phase is dispersed as microscopic droplets in another liquid continuous phase (Desai and Banat, 1997). Poor emulsification of some of the hydrocarbons might be due to the inability of the biosurfactant to stabilize the microscopic droplets. Emulsifying biosurfactants that are stable in environments with high pH and salinity would find applications for bioremediation of spills at seas. The biosurfactant may also be useful for bioremediation works in hot and slightly alkaline environments.

Surface tension measurement

Surface tension is a measurement of the cohesive energy present at an interface. The molecules of a liquid attract each other. The interactions of a molecule in the bulk of a liquid are balanced by an equal attractive force in all directions.

In the present study, strains of *Aeromonas* A1, A2, A3, A4, A5, A6 lowered the surface tension of water to 30, 25, 28, 22, 20, 25, 20mN/m respectively (Table 5). Several strains of anaerobic bacteria produce biosurfactants (Gruła et al., 1983). However, the observed reduction in surface tension (45 to 50 mN/m) was not as large as the observed reduction in surface tension by anaerobic organisms (27 to 50 mN/m) (Cooper, 1986). Lowering of surface tension is an important property of hydrocarbon degrading strains which helps in utilization of the hydrophobic substrates (Rahman, 1993) reported a surface tension value of 29.5mN/m with 1% inoculum at the stationary phase. The glycerol medium with 10% inoculum produced the lowest surface tension value of 29.5mN/m. An effective microbially produced surfactant can lower this value to <30dyn/cm (Lang and Wanger, 1987). The lowest value of surface tension by *Bacillus* sp. and *Aeromonas* sp. was reported by (Takeyama and Matsunaga, 2002). In a study by Oberbremer and Muller-Hurtig (1989), a positive correlation was obtained between reduction in surface tension of the fluid phase in a stirred soil bioreactor and the onset of biodegradation of hydrophobic petroleum hydrocarbons. It has also been previously reported that rhamnolipid can mediate reduction in surface tension (Banat et al., 2000; Noordman et al., 2000).

Biosurfactant estimation


Biosurfactants, amphiphilic compounds of microbial origin, have advantages over their chemical counterparts in biodegradability and effectiveness at extreme temperature, pH and in having lower toxicity (Banat et al., 2000).

In the present study out of 6 strains of *Aeromonas*, A1 shows 0.041g/l of biosurfactant production after 24hrs and at 168 hrs 0.064g/l, A2 shows 0.049g/l at 24hrs and at 168 hrs 0.067g/l, A3 shows 0.043g/l at 24hrs and at 168 hrs 0.064g/l, A4 shows 0.038g/l at 24hrs and at 168 hrs 0.056g/l, A5 shows 0.052g/l at 24hrs and at 168 hrs 0.067g/l, A6 shows 0.034g/l at 24hrs and at 168 hrs 0.049g/l respectively among which the maximum biosurfactant production was show by the A1 at 168hours (Table 6 and Figure 4).

Biosurfactants have been reported to be produced on water soluble compounds such as glucose, sucrose, glycerol, or ethanol (Desai and Banat, 1997). Biosurfactant produced from water-soluble substrates have been reported to be inferior to that obtained with water immiscible substrates (Syldatk et al., 1985; Robert et al., 1989). Such biosurfactants may however be cheaper to produce and useful in food and pharmaceutical industries as it will not required extensive purification. Biosurfactants producing microorganisms may play an important role in the accelerated bioremediation of hydrocarbon contaminated sites (Banat et al., 2000; Rosenberg et al., 1999). Most microbial surfactants are substrate specific, solubilizing or emulsifying different hydrocarbons at different rates (Ilori and Amund, 2001). Moreover, use of biosurfactant producing, hydrocarbon degrading, microorganisms for bioaugmentation to enhance hydrocarbon degradation offer the advantage of a continuous supply of a non-toxic and biodegradable surfactant at a low cost (Moran et al., 2000; Rahman et al., 2000c). Environmental factors such as pH, salinity and temperature also affects biosurfactants activity.

Our results indicate that *Aeromonas* sp. were efficient in biosurfactant production and hydrocarbon emulsification. These results suggests that the use of carbon source like diesel for the biosurfactant production, could enhance the biosurfactant production and there by these strains may be applied for bioremediation of hydrocarbon-contaminated sites and enhanced oil recovery.

4. Figures and Tables







Table 1. Growth range determination of *Aeromonas*

STRAINS	TURBIDITY
Control	-
A1	++
A2	++
A3	++
A4	+
A5	++
A6	+

Table 2. Degradation of diesel by *Aeromonas*

Incubation Periods (Days)	Strains /O.D Value						
	Control	A1	A2	A3	A4	A5	A6
0	0.000	0.251	0.308	0.203	0.111	0.184	0.114
1	0.000	0.257	0.313	0.212	0.117	0.193	0.124
2	0.000	0.264	0.319	0.219	0.121	0.198	0.127
3	0.000	0.271	0.324	0.222	0.129	0.207	0.131
4	0.000	0.279	0.329	0.227	0.133	0.216	0.136
5	0.000	0.284	0.334	0.231	0.139	0.223	0.141
6	0.000	0.289	0.339	0.237	0.143	0.229	0.146
7	0.000	0.293	0.342	0.241	0.147	0.234	0.149

Table 3. Biomass Production by Aeromonas

Incubation Periods (Days)	Strains/ Biomass					
	A1	A2	A3	A4	A5	A6
2	0.05	0.03	0.05	0.02	0.03	0.02
4	0.07	0.05	0.06	0.04	0.04	0.03
6	0.09	0.06	0.08	0.05	0.06	0.04

Table 4. Emulsification activity by Aeromonas

Strains	Hydrocarbon	Percent Emulsified
A1	Diesel	75
A2	Diesel	57.5
A3	Diesel	63.75
A4	Diesel	25
A5	Diesel	35
A6	Diesel	18

Table 5. Surface Tension Measurement

Strains	Surface tension (mN/m)
A1	30
A2	25
A3	28
A4	22
A5	24
A6	21

Table 6. Estimation of biosurfactant by Aeromonas

Incubation Periods (Days)	Strains/OD Values						
	Control	A1	A2	A3	A4	A5	A6
1	0.000	0.041	0.049	0.043	0.038	0.052	0.034
2	0.000	0.047	0.053	0.047	0.041	0.054	0.036
3	0.000	0.053	0.057	0.051	0.044	0.058	0.039
4	0.000	0.057	0.060	0.054	0.047	0.060	0.041
5	0.000	0.059	0.062	0.057	0.051	0.062	0.044
6	0.000	0.061	0.064	0.061	0.054	0.064	0.047
7	0.000	0.064	0.067	0.064	0.056	0.067	0.049

IV. CONCLUSION

The discovery of petroleum brought a lot of relief to the world's energy requirement because of ease of sourcing and conversion. The ease of production, refining and distribution has also brought with it an ever-increasing problem of environmental pollution. One of the ways through which petroleum pollutants can be removed is by solubilization and emulsification. Hydrocarbon oxidizing bacteria, fungi and algae are distributed widely in nature. Fertile soil contains significant number of microorganisms that can utilize hydrocarbon as sole source of carbon and energy. As a hydrocarbon mixture, diesel is obtained in the fractional distillation of crude oil between 250⁰C and 350⁰C at atmospheric pressure. It is generally simpler to refine than gasoline. The purpose of present study was to enumerate the diesel degrading *Aeromonas* and its emulsification activity and production of biosurfactant. The organism used in this study was isolated from oil contaminated soil, the hydrocarbon substrate specificity test showed that diesel is also one of the best substrate for growth and emulsification of biosurfactant by *Aeromonas*. Among six strains of *Aeromonas*, first strain shows maximum degradation rate and emulsification of biosurfactant. About 75% of diesel was degraded by *Aeromonas* over a period of 7 days. The biosurfactant produced by the diesel degrading *Aeromonas* sp. represents a new type of biosurfactant with strong emulsifying ability.

V. FUTURE PROSPECTS

Identification of *Aeromonas* strains with proper identity and identification of other microbes which are inhabited in the same natural environment will give a clear idea about the microbial diversity in the particular automobile service station environment – where petroleum hydrocarbon and detergent levels would be very high. Biosurfactant productivity studies, identification and characterization of biosurfactants by TLC and HPLC will be highly applicable to understand and interaction between microbes and their environment. Emulsification and hydrocarbon degradation studies in the future will give a clear picture about the degradation of petroleum compounds by *Aeromonas* sp.

ACKNOWLEDGEMENTS

“The gracious deeds of the **ALMIGHTY** are beyond the words of compare”. With a warm heart, I thank God for his blessing showered upon me to complete my project work.

I accord my deep sense of gratitude to the founder of K.S.Rangasamy, Educational Institutions, **LION.Dr.K.S.RANGASAMY, MJF.**, Executive Directress of K.S.Rangasamy College of Arts and Science,

I express my gratitude and sincere thanks to **Dr. M.GANESAN**, Principal, K.S.R College of Arts and Science and **Dr. N. KANNAN**, Hon. Reader, Department of Biotechnology for their encouragement ,

I owe my true and profound sense of gratitude to thank **Dr. G. GURUSUBRAMANIAN, M.Sc., D.Ent., Ph.D., D.T.M.E., D.C.A.**, PG and Research Department of Microbiology, for his valuable suggestion and successful completion of my project.

I express my whole hearted thanks to **Dr. G. VIVEKANANDHAN, M.Sc., M.Phil., Ph.D., Lecturer**, PG and Research Department of Microbiology, K.S.R College of Arts and Science for his keen interest, meticulous support, valuable, enthusiastic guidance and brilliant suggestions throughout my project work.

I also avail this opportunity to thank to **Dr. A. SANKARANARAYANAN, M.Sc., Ph.D.**, Head, PG and research department of Microbiology, for his valuable support and assistance that were readily offered to me throughout the course of my study.

I express my whole hearted thank to **Dr. K.S.M.Rahman, Biotechnology Research Group, School of biological and environmental sciences, University of Ulster, Coleraine, country Londonderry, Northern Ireland.BT 521SA UK** for his valuable support and assistance that were readily offered to me throughout the course of my study.

REFERENCES

- [1]. Alexander. M. 1995. Biodegradation and bioremediation Chapter 2 New York Academic press.
- [2]. Amikam Harounitz., David Gutnick and Eugene Roserberg. 1975. Sequential growth of bacteria on crude oil. Applied microbiology, 30: 10 – 19.
- [3]. Apalijalahti. JHA. and Salkinoja Salonen M.S. 1984. Absorption pentachlorophenol (PCP) by bark chips and its role microbial PCP degradation. Microb. E.coli.10: 359 – 367.
- [4]. Atlas. R. M. and R. Bartha. 1972. Degradation and mineralization of petroleum by two bacteria isolated from coastal waters. Biotechnol. Bioeng. 14:297-308.
- [5]. Atlas. R.M. 1981. Fate of oil from two major oil spills. Environ. It 5: 33 – 38.,Microbial degradation of petroleum hydrocarbons, an environmental perspective. Microbial Rev. 45: 180 – 209.
- [6]. Atlas. R.M. 1990. Microbial hydrocarbon degradation-bioremediation of oil spills. J. Chem. Tech. Biotechnol. 52:149-156.
- [7]. Atlas. R.M. 1995. Bioremediation of petroleum pollutants. Int. Biodeterior. Biodegrad., 21, 317-327.
- [8]. Atlas. R.M. and Bartha. R. 1992. Hydrocarbon biodegradation and oilspill bioremediation Adv. Microb. Eco. 12:287.338.
- [9]. Atlas. R.M. and R. Bartha. 1972. Biodegradation of petroleum in sea water at low temperatures. Can. J.Microbiol. 18:185–195.
- [10]. Banat. I. M., R.S. Makkar and S.S. Cameotra. 2000. Potential commercial application of microbial surfactants. Appl. Microbiol. Biotechnol. 53: 495-508.
- [11]. Banat. I.M. 1995. Biosurfactants production and use in microbial enhanced oil recovery and pollution remediation. A review. Biores. Technol. 51, 1-12.
- [12]. Banerjee. S., Duttagupta, S. and Chakrabarty, A.m., 1983. Arch. Microbiol. 135, 110 – 114
- [13]. Barathi. S., and Vasudevan. N. 2001. Utilization of petroleum hydrocarbons by *Pseudomonas fluorescens* isolated from a petroleum – contaminated soil. Env. Int., 26:413-416.
- [14]. Bartha. R. 1986. Biotechnology of petroleum pollutant Biodegradation microbial Ecol. 12: 155 – 172.
- [15]. Bauchop. T., and Elsdon. S.R. 1969. The growth of microorganisms in relation to their energy. J. Gen. Microbiol. 23,457-469.
- [16]. Berg. G., Seech, A.G. Lee, H. and Trevors, j. T., 1990. J. Environ. Sci. Health. A 25, 753 – 764.
- [17]. Bergey. D.H. 1939. Manual of Determinative Bacteriology. 5th Ed., Baltimore, Williams and Wilkins Company, 1032pp.
- [18]. Blumer. M. 1971. Scientific aspects of the oil spill problem. Environ. Aff. 1:57-73.
- [19]. Bossert. I. and Bartha. R. 1984. The fate of petroleum and soil ecosystems. In: Petroleum Microbiology, ed. Atlas, R.M. New York: Macmillan Publishing Co. 434-476.
- [20]. Broderick. L.S. and Cooney. J.J. 1982. Emulsification of hydrocarbons by bacteria from fresh water ecosystems. Dev.Ind.microbiol. 23: 425 – 434.
- [21]. Candler. JE., Rushing. JH. and Leuteman. AJJ. 1993. Synthetic-based mud systems. SPE Paper 25993, In: SPE/EPA. Exploration and Production Environmental Conference, San Antonio, Texas, USA, 7-10 March 485-499.
- [22]. Chakrabarty. A.M. 1976. Plasmids in *Pseudomonas*. Annu.Rev.Genet. 10: 7 – 30.
- [23]. Chhatre. S., Purohit. H.J., Shanker. R and Khanna. P. 1996. Bacterial consortia for crude oil spill remediation. Water Sci. Technol. 34, 187-193.

- [24]. Conomoe. T.J. 1975. Movement of applied oil an predicted by estuarism nontidal drift. *Limnol. Oceanogr.* 20: 159-173.
- [25]. Del' Arco, J.P and De Franca. F.P. 1999. Biodegradation of crude oil in sendy sediment. *Int. Biodeter. Biodegrad.* 44, 87-92.
- [26]. Desai, A. J., Patel, K.M. and Desai, J.D. 1987,88. *Curr. Sci.* 57, 500-501, Microbial surfactants evaluation, tyupes and future applications placations. *J.Sci.Ind.Res.* 46: 440 – 449.
- [27]. Desai, J.D. and Banat. I.M. 1997. Microbial production of biosurfactants and their commercial potential. *Microbiol. Mol. Bio. Rev.* 61: 47 – 64.
- [28]. Dibbel. J.T. and R. Bartha. 1976. Leaching aspects of oil sludge biodegradation in soil. *Sci.*, 127 : 365 – 370.
- [29]. Donaldson. E.C., G.V. Chilingarian and T.F. Yen. 1989. Introduction. In *Microbial enhanced oil recovery*. Elsevier, New York, N.Y. 1-15.
- [30]. Duvnjak. Z., Cooper. D.G., and Kosaric. N. 1983. Effect of nitrogen source on surfactant production by *Arthrobacter paraffines* ATCC 19558. In: Zajic, J.E., Cooper, D.G., Jack, T.R., Kosaric, N. (Eds.), *Microbial Enhanced Oil recovery*. Pennwell Books, Tulsa, Oklahoma, USA, 66-72.
- [31]. Ellis. W.D., Payne, J.R. and Monabb G.D., 1985. Treatment of Contaminatn soils with Aqueous Surfactants, United States environmental Protection Agency, Office of Research and Development.
- [32]. Enrlieh. G.G., Schroeder. R.A. and Martin. P., 1985. Microbial Populations in a jet Fuel contaminated Aquifer at Tustin, California, US geological Survey Report, 85.
- [33]. Fehler. S.W.G and R.J. Light. 1970. Biosynthesis of hydrocarbons in *Anabaena variabilis*. Incorporation of (methyl-C) and (methylH) – methionine into 7-and 8-methyl-heptadecanes. *Biochemistry.* 9:418-422.
- [34]. Ferrari. M.D., Neirotti. E., Albornoz. C., Mostazo. and M.R., Cozzo. M. 1996. Biotreatment of hydrocarbons from petroleum tank bottom sludges in soil slurries. *J. Biotechnol. Lett.*, 18:1241-1246.
- [35]. Fingas. M, Environment Canada, Personal communication, 1990.
- [36]. Floodgate. G. 1984. The fate of petroleum in marine ecosystems. In: *Petroleum Microbiology*, ed. Atlas, R.M. New York: Macmillan Publishing Co. 335-398.
- [37]. Fought. J. M., Gutnick. D.L and Westlake. D.W.S., 1989. *Appl. Environ. Microbiol*, 55, 36 – 42.
- [38]. Frankenberger. Jr. W.T. 1992. The need for a laboratory feasibility study in bioremediation of petroleum hydrocarbons. In: Calabrese, E.J.; Kostecki, P.T. (eds.) *Hydrocarbon contaminated soils and groundwater*. Lewis Pub., Boca Raton, FL, 237-293.
- [39]. Gary Canny (Email: cannyg@students. Itsligo i.e., Dr. Michael Broaders (Email. broaders, Michael@itsligo.ie., 1997.
- [40]. Gatellier. C.R. 1971. Less features limitant la biodegradation deshydrocarbures dans puration des eaux. *Chim.Ind (Paris)* 104: 2233 – 2289
- [41]. Gillam. A.H., and Gibson. M.J. 1986. Chemical monitoring in the North sea. *Chemistry in Britain.* Oct. 911-913.
- [42]. Grula. E.A., Russell. H.H., Bryant. D., Kenaga. M and Hart. M., in Proceedings of the 1983. International conference on microbial department of Energy, Bartlesville, 43.
- [43]. Gunkel. W. 1968. Bacteriological investigations of oil polluted sediments from the Cornish coasts following the “Torrey canyon” disaster. In J.D Carthy and D.R Arthu (Ed). *The biological effects of oil pollution on littoral communities (Supplement to field Stud. 2)* E.W.classsey, Ltd.,
- [44]. Hardwood. J. L. and N.J. Russel. 1984. *Lipids in plants and microbes*, george Allen and Unwin Ltd. London, U.K. 110-111.
- [45]. Hommel. R.F. and C. Ratledge. 1993. Biosynthetic mechanisms of low molecular weight surfactants and their precursor molecules. 3 – 63.
- [46]. Hommel. R.K. 1990. Formation and physiological role of biosurfactants produced by hydrocarbon-utilizing microorganisms. *Biodegradation* 1:107-110.
- [47]. Hunt. J. M., R.J. miller and J.K. Whelan. 1980. Formation of C-C hydrocarbons from bacterial degradation of naturally occurring terpenoids. *Nature (London)*, 288:577-578.
- [48]. Ilori. M.O., and Amund. O.O. 2001. Production of a peptidoglycolipid bioemulsifier by *Pseudomonas aeruginosa* grown on hydrocarbon. *Z. Naturforsch.* 56C, 547-552.
- [49]. Ilori. M.O., C.J. Amobi and A.C. Odacha. 2005. Factors affecting biosurfactant production by oil degrading *Aeromonas* spp. isolated from a tropical environment. *Chemosphere* 61: 985 – 992.
- [50]. Iqbal. S., Khalid. Z.M. and Malik K.A. 1995. Enhanced biodegradataion and emulsification of crude oil and hyper production of biosurfactants by a gamma ray induced mutant of *Pseudomonas aeruginosa*. *Let Appl. Micro.* 21: 176 – 179.
- [51]. J.D. Walker., R.R. Colwell and L. Petrakis. 1975. Evaluation of petroleum degrading potential of bacteria from water and sediments. *Applied microbiology* 30 (6), 1036 – 1039.
- [52]. Jain. D.K., Lee. H and Trevors. J.T. 1992. Effect of addition of *P. aeruginosa* UG2 inocula or biosurfactants on biodegradation of selected hydrocarbons in soil. *J. Ind. Microbiol.* 10:87-93.
- [53]. Jobson. A., F. D. Cook and W.S. Westlake. 1972. Microbial utilization of crude oil. *Appl. Microbiol.* 24:1082-1088.
- [54]. Jobson. A., M. McLaughlin, F.D. Cook and D.W.S. Westlake. 1973. Effect of Amendments on the microbial utilization of oil applied to soil. *Applied. Microbiology.* 166-171.
- [55]. John Wiley and Sons. 1982. *Encyclopedia of chemical Technology* 3rd ed. 17:119-130.
- [56]. Juttner. F. 1976. β -Cyclocitral and alkanes in microcystis (Cyanophyceae). *Zeitschrift fur Naturforschung.* 31c, 491-495-503.
- [57]. Kaja Sheik Mujibur Rahman, Thanhira Rahman, Perumal Lakshmanaperumalsamy and Ibrahim M. Bana. 2003. Occurrence of crude oil degrading bacteria in gasoline and diesel station soils. *J.Basic.Microbial.* 42: 286 – 293.
- [58]. Karanth, N.G.K., P.G. Deo and N.K. Veenanadig. 1999. Microbial production of biosurfactants and their impotence.
- [59]. Kaserer. H. 1906 Uber die Oxydation des Waserstoffes and des Mothans durch Mikroorganismen. *Zentr. Bakt. Parasitenk., Abt II*, 15: 573-576.
- [60]. Kastner. M., Breuer-Jammali, M and Mahro. B. 1994. Enumeration and characterization of the soil microflora from hydrocarbon-contaminated soil sites able to mineralize polycyclic aromatic hydrocarbons (PAH). *Appl. Microbiol. Biotechnol.* 41, 267-273.
- [61]. Kastner. M., Breuer-Jammali and M., Mahro, B. 1994. Enumeration and characterization of the soil microflora from hydrocarbon-contaminated soil sites able to mineralize polycyclic aromatic hydrocarbons (PAH). *Appl. Microbiol. Biotechnol.* 41, 267-273.
- [62]. Klevans. H.B. 1954. Solubilization of polycylic hydrocarbons *J. Phys. Colloid Chem.*, 54: 283-298.
- [63]. Kolattukudy. P.E., J.S. Buckner and L. Brown. 1972. Direct evidence for a decarboxylation mechanism in the biosynthesis of alkanes in *B. oleracea*. *Biochem. Biophys. Res. Commun.* 47:1306-1313.
- [64]. Kornelli. T.V., Kouroua, T.L. and Yu. V. Denisov. 1983. *Microbiology* 52, 767-770.
- [65]. Kosaric. N., Cairn. W.L.S., Gray and N.C.C. (eds) 1987. *Biosurfactants and Biotechnology*, surfactant Science Series, vol. 25, Marcel Dekker.

- [66]. Kretschmer. A., Bock. H and Wagmer. F. 1964. Characterisation of biosynthetic intermediates of trehalose dicorynemycolates from *Rhodococcus erythropolis* on n-alkanes, *Appl. Environ. Microbiol.*, 44 (1982).
- [67]. Kroos. H. domestic and International, Atlanta, 1989.
- [68]. Lal. B and Khanna. S. 1996. Degradation of crude oil by *Acinetobacter calcoaceticus* and *Alcaligenes odorans*. *J. Appl. Bacteriol.* 81, 355-362.
- [69]. Lang S., and F. wagner. 1987. Structure and properties of biosurfactants, p.,21 – 47. In N.Kosaric, W.L Cairns, and N.C.C. Gray (ed.), *Biosurfactants and biotechnology*. Marce Deker, Inc., new York, N.Y.
- [70]. Lang. S and Wanger. F. 1987. Structure and properties of biosurfactants. *Surfactant Science Ser* 25: 21-45.
- [71]. Leahy. J.G. and R.R. Colwell. 1990. Microbial degradation of hydrocarbons in the environment. *Microbiol. Rev.* 54:305-315.
- [72]. Margesin. R. and F. Schinner. 1997. Efficiency of Indigenous and Inoculated cold-adapted soil Micro-organisms for Biodegradation of Diesel Oil in Alpine soils. *Appl. Environ. Microbiol* 67(7): 2660-2664.
- [73]. McClay. K., B.G. Fox and R.J. Steffan. 2000. Toluene monooxygenasecatalyzed epoxidation of alkenes. *Appl. Environ. Microbiol.* 66: 1877-1882.
- [74]. McLnerney. M.J., Javaheri. M and Nagle. D.P. 1990. Properties of the biosurfactant produced by *Bacillus licheniformis* strain JF-2. *J. Ind. Microbiol.* 5, 95-102.
- [75]. Meredith. W., Kelland. S.J and Jones. D.M. 2000. Influence of biodegradation on crude oil acidity and carboxylic acid composition. *Org. Geochem.* 31, 1059-1073.
- [76]. Midwest Biofuels Inc. 1993. Soy Diesel in biodegradability and toxicity technical information. In, *Biodiesel information kit* received from National Soy Diesel Development Board, Overland Park, K.S.
- [77]. Miget. R.J., C.H. Oppenheimer, H.I. Kater and P.A. LaRock. 1969. Microbila degradation of normal paraffin hydrocarbons in crude oil. In proceedings joint conference on prevention na dcontrol of oil spills. API/FWPCA Conference, U.S. Department of Interior, Washington, D.C. 327-331.
- [78]. Mikkelson. J.D., and P. Von Wettstein-Knowles. 1978. Biosynthesis of β -diketones and hydrocarbons in barley spike epicuticular wax. *Arch. Biochem. Biophys.* 188:172-181.
- [79]. Moran. A.C., Olivera. N., Commendatore. M., Esteves. J.L and Sineriz. F. 2000. Enhancement of hydrocarbon waste biodegradation by addition of a biosurfactant from *Bacillus subtilis* 09. *Biodegradation* 11, 65-71.
- [80]. Morgan. P and Watkinson. R.J, 1989. *CRC Crit. Rev. Biotechnol.*, 1989, 8, 305 - 333.
- [81]. Oberbremer. A., Muller Hurtig. R and Wagner. F., 1990. *Appl. Microbial. Biotechnol.* 32, 485 – 489.
- [82]. Obire. O. and I.V. Okudo. 1997. Effects of Crude oil on a Freshwater Stream in Nigeria. *Discov. Innov.* 9:25-32.
- [83]. Obire. O. and Nwaubeta. O. 2001. Biodegradation of Rejined petroleum hydrocarbons in soil. *Journaly of applied sciences, and environmental management.* 5(1) 43 – 46.
- [84]. Okerentugba. P.O and O.U. Ezeronye. 2003. Petroleum degrading potentials of single and mixed microbial cultures isolated from rivers and refinery effluent in Nigeria. *African Journal of Biotechnology* 2(9), 288-292.
- [85]. Olivera. N.L., Esteves. J.L., and Commendatore. M.G. 1997. Alkane biodegradation by a microbial community from contaminated sediments in Patagonia, Argentina. *Int. Biodeter. Biodegrad.* 40, 75-79.
- [86]. Osenga. M. 1993. What exactly is biodiesel? Diesel progress engines and drives. In, *National Soy Diesel Development Board Alternative Fuels Special Report*.
- [87]. Payne. J.R. and McNabb. D.G.Jr. 1984. Weathering of petroleum in the marine environment, *MTS Journal.* 18: 24 – 42.
- [88]. Portier. R., Bianchini., Fujusaki. K., Henery. C. and Macmillin. 1998. Comparison of effective toxicant biotransformation by autochthonous microorganisms and commercially available cultures in the insitu reclamation of abandoned industrial sites, *Schriftner, ver Wasser baden Iuftthy g, Berlin dehlem.* 80 – 273.
- [89]. Prince. R.C. R. Varadaraj, R.J. Fiocco and R. R. Lessard. 1999. Bioremediation as an oil response tool. *Environ. Technol.* 20:891-896.
- [90]. Rahman. K.S.M, I.M. Banat., J. Thahira, Tha. Thayumanavan and P. Lakshmanaparumalsamy. 2002. Bioremediation of garoline contaminated soil by a bacterial consortium amended with poultry litter, coil pith and rhamnidiol Biosurfactant *Bioresource technology.* 81: 25 – 32.
- [91]. Rahman. K.S.M., Thahira-Rahman. J., Lakshmanaperumalsamy. P. and Banat. I.M. 2002c. Towards efficient crude oil degradation by mixed bacterial consortium. *Bioresour. Technol.* 85, 257-261.
- [92]. Ramana, K.V. and Karanth, N.G.K., 1989. *J. Chem. Technol. Biotechnol.*, 45, 249 – 257.
- [93]. Raymond. R.L., Hudson. J.O and Jamison. V.W. 1976. Oil degradation in soil. *J. Appl. Environ. Mcirobiol.* 31: 522 – 35. .
- [94]. Rosenberg. E. 1991. Hydrocarbon Oxidizing bacteria. In the prokaryotes, ed. A.Ballows. 441 – 59.
- [95]. Rosenberg. E., and Ron. E.Z. 1999. High molecular mass microbial surfactants, *J. Appl., Micrbiol, Biotechnol,* 52: 154 – 162.
- [96]. Rosenberg. E., Zukerberg. A., Rubinovitz. C and Gutnick. D.L. 1979. Emulsifier of *Arthrobacter* RAG-1, Isolation and emulsifying properties. *Appl. Environ. Microbial.* 37, 402-408.
- [97]. Schmidt. S.K. and Alexander. M. 1985. Effects of dissolved organic carbon and second substrates on biodegradation of organic compounds at low concentration *J. Appl. Environ, Microbiol.*, 49: 822 – 827.
- [98]. Seklemova. E., Pavlova. A. and Kovacheva. K. 2001. Biostimulation-based bioremediation of diesel fuel: Field demonstration. *Biodegrad.* 12:311-316.
- [99]. Shimp. R.J. and Pfaender. F.K. 1985. Influence of easily degradable naturally occurring carbon substrates on biodegradation of monosubstituted phenol by aquatic bacteria. *J.appl. Environ. Microbial.* 49: 394 – 401.
- [100]. Singer. M.E. and W.R. Finnerty. 1984. Microbial metabolism of straight-chain and branched alkanes. In, R.M. Atlas (ed.), *Petroleum Microbiology*, Macmillan Publishing, New York. 1-60.
- [101]. Sohngen. N.L. 1906. Uber Bakterien, welche Methan als Kohlenstoffnahrung Energiquelle gebrauchen. *Zentr. Bakt. Parasitenk., Abt. II,* 15; 513-517.
- [102]. Steffan. R. J., K. Mcclloy., S. Vain berg., C.W. Condee and D. Zhang, 1997. Biodegradation of the Gasoline Oxygenates Methylene tert-Butyl Ether, Ethyl tert-Butyl Ether and tert-Amyl Methyl Ether by Propane-Oxidizign Bacteria. *Appl. Environ. Microbiol.* 63(11): 4216-4222.
- [103]. Stella. O., Ajisebutu. 1988. Effectsof sodium chloride on biodegradation of crude oils by two species of *Aeromonas*. *Applied. Microbiology and Biotechnology.* 28 (2), 203-208.
- [104]. Stone. R.W., White. A.G.C. And Fenske. M.R. 1940. Microorganisms attacking petroleum and petroleum factions. *lbact.*, 39: 91 – 92.
- [105]. Sugiura. K., Ishihara. M., Shimauchi. T and Harayama. S. 1997. Physicochemical properties and biodegradability of crude oil. *Environ. Sci. Technol.* 31, 45-51.

- [106]. Syldatk. C. and F. Wagner. 1987. Production of biosurfactants. In n.Kosaric, W.L.Cairns, N.C.C gray, (ed) Biourfactants and biotechnology Marcel Dekker, Newyork. 89 – 12.
- [107]. Takeyama. H., Wada. M. and Masunaga. T. 2002. Screening of soil bacteria for production of biocleaner. *Applied Biochemistry and Biotechnology*, 98 (1-3), 319-326(8).
- [108]. Traxler. R.W. 1973. Bacterial degradation of petroleum materials in low temperature marine environments. 163 – 170.
- [109]. Van Dyke. M.L., H. Lee and J.T. Trevers. 1991. application of microbial surfactants. *Adv.*, 9. 241-252.
- [110]. Vasudevan. N. and Rajaram. P. 2001. Bioremediation of oil sludge-contaminated soil. *Environ. Int.* 26, 409-411.
- [111]. Venkateswaran. K. and Harayama. S. 1995. Sequential enrichment of microbial populations exhibiting enhanced biodegradation of crude oil. *Can. J. Microbiol.*, 41, 767-775.
- [112]. Vivekanandhan. G., N. Vasudevan and P. Lakshmanaperumalsamy. 1999. Degradation of crude oil by a bacterial consortium. 18(3), 245-249.
- [113]. Voroshilova. A. A. and E.V. Disnova. 1950. Bacterial oxidation of oil and its migration in natural waters *Mikrobiologiya* 19:203-210.
- [114]. Walker, J.D and Colwell, R.R. 1974. Microbial degradation of model petroleum at low temperatures. 1, 63-95.
- [115]. Walker. J.D., R.R. Colwell. 1976a. Measuring potential activity of hydrocarbon degrading bacteria. *Appl. Environ. Microbiol.* 31:189-+197.
- [116]. Walker. J.D., and R. Colwell. 1976b. Enumeration of petroleum-degrading microorganisms. *Appl. Environ. Microbiol.* 31:198-207.
- [117]. Ward. D.M., R.M. Atlas, P.D. Boehm and J.A. Calder, 1980. "Microbial Biodegradation and the Chemical Evolution of Amoco codiz Oil Pollutants" *Ambio*, vol. IX, No. 6, 279.
- [118]. Wasko. M.P and Bratt. R.P. 1990. Properties of a biosurfactant produced by the fuel contaminant *Ochrobactrum anthrppii*. *Inter. Biodeter.*, 27, 265 – 73.
- [119]. Wilkinson, S.G and Cordell. 1985 Studies on lipopolysaccharides from *Pseudomonas aeruginosa*. *Eur. J.Biochem.* 52: 331 – 343.
- [120]. Winters. K., P.L. Parker and C. Van Baalen. 1969. Hydrocarbons of the blue-green algae: Geochemical significance. *Science.* 163:467-468.
- [121]. Wong. J.H.C., Lim. C.H. and Nolen. G.L. 1997. *Design of Remediation Systems*; Lewis Publishers Ed.; United States of America.
- [122]. Yeung. P.Y., Johnson. R.L., Xu. J.G. 1997. Biodegradation of petroleum hydrocarbons in soil as affected by heating and forced aeration. *J. Environ. Qual.* 26, 1511-1516.
- [123]. Zajie. J.E., Guignard. H. and Gerson. D.F, 1977. Properties of biodegradation of bioemulsifier from *corynebacterium hydroclastus*, *Biotechnol Bioeng.* 191-303.
- [124]. Zhang. Y., and Miller. R.M. 1992. Enhanced octadecane dispersion and biodegradation by a *Pseudomonas rhamnolipid* surfactant (biosurfactant). *Appl. Environ. Microbial.* 58, 3276-3282.
- [125]. ZoBell. C. E. and J.F. Prokop. 1946. Microbial oxidation of mineral oils in Barataria Bay bottom deposits. *Z. Allg. Mikrobiol.* 6:143-162.
- [126]. ZoBell. C.E. 1969. Microbial modification of crude oil in the sea, p.371-326. *Proc. API/FWPCA Conference on Prevention and control of Oil Spills.* Atherican Petroleum Institute publ. No.4040.