

Evaluation of Toxic Effect of Terminalia Chebula Fruit Extract in Albino Rats

Elavarasi, S., Horne Iona Averal, Nevika, E And P. Kanimozhi

PG and Research Department of Zoology, Holy Cross College (Autonomous), Tiruchirappalli.

* Corresponding Author : Elavarasi, S.

ABSTRACT: *The present study aims to analyze the phytochemicals and test the toxic effect of the fruit extract of Terminalia chebula using three different solvents via ethanol, acetone and benzene by examining the changes in behavior, body weight, food intake, water intake, hematological parameters (total WBC count, WBC differential count, RBC, Hb, HCT, MCV, MCH, MCHC and platelet count) and histological changes in the vital organs such as lungs, heart, liver and kidney. No behavioral changes or any toxic symptoms and mortality were observed throughout the experimental period. There is a slight increase or decrease in the body weight of the extract treated groups compared to control group rats. The food intake and water intake of the control and extract treated rats showed slight variations, but not showed the abnormal intake of food and water throughout the experimental period. The hematological parameters showed significant difference among the different extract treated rats and control rats, but the levels are not exceeded from the normal range. The microscopic and macroscopic examination of the vital organs such as lungs, heart, liver and kidney showed normal cell structures, blood vessels and nuclei. Thus the present study revealed that the ethanol, acetone and benzene extracts of Terminalia chebula fruit did not produce any toxic effects of the high dose of 2000mg/kg body weight and is found to be safe. Thus, it is concluded that the plant extract of Terminalia chebula fruit up to 2000 mg/kg body weight was used for further evaluation studies.*

KEY WORDS: *Terminalia chebula, fruit extract, phytochemical analysis, Rattus norvegicus. toxicity studies*

Date of Submission: 19-04-2018

Date of acceptance: 04-05-2018

I INTRODUCTION

India is richly endowed with a wide variety of plants having medicinal value. These plants are widely used by all sections of the society either directly as folk remedies or indirectly as a pharmaceutical preparation of modern medicine (Rajesh et al., 2010). Plants have formed the basis of sophisticated traditional medicine systems that have been in existence for thousands of years (Mahidol et al., 2003). Medicinal plants are plants that have at least one of their parts (leaves, stem, barks or roots) used for therapeutic purposes (Bruneton, 1993). Recently, medicinal plants have become important for the treatment of different disease conditions, such as diabetes, malaria, anaemia (Fola, 1993). The availability and relatively cheaper cost of medicinal plants, makes them more attractive as therapeutic agents when compared to ‘modern’ medicines (Agbor and Ngogang, 2005). Phytochemicals with biological activity have had great utility as pharmaceuticals and pharmacological actions (Rajesh et al., 2010). In recent years, because of their easy availability and cost effectiveness, there has been a dramatic rise in use of herbal drugs/preparations in the developed countries besides having desired pharmacological effectiveness with high level of safety and low toxicity profile (Kadam et al., 2012).

Plants have been used in medicines since time immemorial. India has a rich heritage of using medicinal plants in traditional medicines, as in the Ayurveda, Siddha and Unani systems besides folklore practices. The earliest inscription of the medicinal uses of plants is found in the “Rigveda”, which is one of the oldest repositories of human knowledge. Fairly comprehensive information on the curative properties of some herbs has been found recorded in “Charak Samhita” and “Sushruta Samhita” (Kamboj, 2000). The plant kingdom is a virtual goldmine of biologically active compounds and it is estimated that only 10-15% of existing species of higher plants have been surveyed. Many plants have been successfully used in the treatment of various diseases. The ancient record is evidencing their use by Indian, Chinese, Egyptian, Greek, Roman and Syrian dates back to about 5000 years. In India, around 20,000 medicinal plant species have been recorded recently, but more than 500 traditional communities use about 800 plant species for curing different diseases (Kamboj, 2000). Phytochemistry has been instrumental in the rationalization of the use of various herbal medicines, however unscreened herbal products still find their way to markets owing to their high demand. The quest to unravel the mysteries of bioactive properties of medicinal plants and the comprehension of their nutritional and toxicological constituents have been a subject of intense renewed interest for many scientists all over the World (Ugbogu et al., 2016).

Phytochemicals come in a variety of forms and different vegetables have higher concentrations of a particular phytochemical than others. Some of the main phytochemicals include: carotenoids, isoflavonoids, steroids, lignans, anthraquinones, gums and resins (Rajesh et al., 2010). The most important of bioactive constituents of plants are alkaloids, tannins, flavonoids and phenolic compounds (Doss, 2009). Toxicological studies help to decide whether a new drug should be adopted for clinical use or not depending on the duration of exposure of animals to drug, toxicological studies may be of three type viz. acute, sub-acute, and chronic. Thus the present study focused on the preliminary phytochemical analysis to know the bio-active compounds of the medicinal plant Terminalia chebula and in order to know bio-safety of the Terminalia chebula, acute toxicity test is to be performed through in vivo studies.

II MATERIALS AND METHODS

The selected plant material for the present study is the fruits of Terminalia chebula. The process of extraction and formulation is as described by Sohini et al., (1996). The fruit powder of Terminalia chebula was pulverized and extracted as a whole preparation in a Soxhlet apparatus using polar (ethanol and acetone) and non-polar (benzene) solvents. The different extracts of Terminalia chebula fruit were concentrated to a dry mass by vacuum evaporator and stored in desiccator. The percentage yield was obtained using this formula $W2-W1/W0 \times 100$ (Where, W2 is the weight of the extract and the container, W1 is the weight of the container alone and W0 is the weight of the initial dried sample).

Preliminary phytochemical screening

The fruit powder of Terminalia chebula was subjected to analyse the preliminary phytochemicals such as alkaloids, carbohydrates, Fixed oils and fats, flavonoids, glycosides, phenolic compounds, protein, steroids, saponins, tannins and terpenoids according to the standard methods (Kokate, 1994; Harborne, 1973; Rajpal, 2002; Raaman, 2006).

Drug dosage calculation is followed by the method of Erhirhie et al., 2014. Experiments on animals are necessary in drugs discovery and development as well as to advance medical and biological knowledge (Baker et al., 1979). Dosage calculation and stock solution preparation based on dosage rationale formula are prerequisites to drug administration in experimental animals (Erhirhie et al., 2014).

Experimental set up for acute toxicity study

Healthy adult male Wistar Albino rats, *Rattus norvegicus* (150-200 mg/kg b.wt.) were used for the present study. The rats were obtained from SASTRA University, Thanjavur and brought to the laboratory and maintained under controlled environment. The rats were randomized into control and experimental groups and housed in different plastic cages. All animals were fed with standard pellet feed and water ad libitum. The principles of animal care (Ethical Committee's Approval No.001/HCC/IAEC/DST-NPDF/2017) were followed throughout the experimental period.

Normal, healthy male albino rats fasted for 12 hours were randomly divided into control and extracted treated groups. There were three different types of extracts (ethanol extract, acetone extract and benzene extract of Terminalia chebula) separately tested for their toxic effect. In each extract, 2000mg/kg was separately administered orally to the rats for 14 days. The rats were observed for clinical signs and symptoms of toxicity and mortality from the time of extract administration to 14th day. Behavioural changes, changes in body weight, daily food intake and water intake were observed over a period of 14 days. At the end of the experiment, all experimental animals were fasted overnight, cervically dislocated, and blood samples were collected by cardiac puncture into tube with anticoagulant ethylene di-amine tetra acetic acid (EDTA) for haematology. Blood samples in test tubes containing EDTA were immediately processed for haematological parameters using Automated Haematological Analyzer. Total and differential white blood cell count (WBC), red blood cell count (RBC), the haemoglobin concentration (HGB), hematocrit (HCT), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), Red cell distribution width, and platelet count (PLC) were determined. And the vital organs such as lungs, liver, kidney and heart tissues were removed and washed with ice cold saline and, weighed and preserved in 10% formalin solution for histological studies.

Statistical analysis:

Values were represented as Mean \pm Standard deviation. All statistical analyses were performed by using windows based SPSS package (Statistical Package for Social Sciences / Statistical Product and Service Solutions).

III RESULTS AND DISCUSSION

Percentage of extraction:

The yield of crude ethanol extract of *T. chebula* fruit (20.76%) is the highest among the three samples whereas yield of a crude acetone extract of *T. chebula* fruit (10.98%) and the crude benzene extract of the *T. chebula* fruit is 7.25%.

Preliminary Phytochemical analysis:

Phytochemical analysis concluded on the plant extracts revealed the presence of constituents which are known to exhibit medicinal as well as physiological activities (Sofowra, 1993). In the present study acute toxic effect of ethanol, acetone and benzene extract of *Terminalia chebula* fruit was evaluated. It revealed the presence of many phytochemical compounds which possessed various medicinal activities. The ethanol extract of *T. chebula* fruit powder revealed the presence of phytochemicals such as protein, phenol, flavonoids, tannin, sterol and carbohydrates. The acetone and benzene extract of *T. chebula* fruit powder showed the presence of protein, phenol, flavonoids, tannin, saponin, terpenoids and carbohydrates. The phenolic compounds are one of the largest and most ubiquitous groups of plants metabolites (Singh et al., 2008). They possess biological properties such as anti-apoptosis, anti-aging, anti-carcinogen, anti-inflammations, cardiovascular protection and the improvement of endothelial function, as well as inhibition of angiogenesis and cell proliferation activities. Several studies have described the antioxidant properties of medicinal plants which are rich in phenolic compounds (Krings and Berger, 2001). Natural antioxidants mainly come from plants in the form of phenolic compounds such as flavonoid, phenolic acids, tocopherols etc., (Ali et al., 2008). Flavonoids have antioxidant activities as well as health promoting effects viz., anti-allergic, anti-cancer, anti-oxidant, anti-inflammatory, anti-thrombotic, vasoprotective, tumor inhibitory and anti-viral effects (Salah et al., 1995; Okwu, 2004; Hodek et al., 2002). Phytochemicals such as saponins, terpenoids, flavonoids, tannins, steroids and alkaloids have anti-inflammatory effects (Akindale and Adeyemi, 2007). Thus the phytochemicals present in the plant extract revealed that the plant drug possess anti-inflammatory, antibacterial, antioxidant, anti-allergic, anticancer, etc., may be used to treat several diseased conditions especially used for skin diseases.

Acute toxic effect of the plant extract on behavioural changes in rats

Herbal medicine preparations are "natural" and are therefore intrinsically harmless. However, their effects can be very powerful and potentially lethal if used incorrectly and their use as a substitute for conventional medicines may be ineffective (Bateman et al., 1998). In the present study, acute toxic effect of ethanol, acetone and benzene extract of *Terminalia chebula* fruit powder was evaluated. There were no noticeable changes in the general behaviour, toxicity signs and mortality observed in rats treated with test drug orally at 2000 mg/kg body weight for a period of 14 days.

Toxic effect on body weight

Weekly body weight changes among the different extract treated rats and control rats are shown in Figure 1. The control rats and the extract treated rats showed a normal increase in their body weight throughout the experimental period. The body weight of control rats are 207.8 ± 8.60 and at the end of the experiment it increased up to 216.3 ± 10.98 g. The ethanol, acetone and benzene extract treated rats showed the body weight of 193.4 ± 8.41 g, 190.4 ± 4.92 g and 190.1 ± 4.15 g, and at the end of experimental period it reached about 195.9 ± 8.67 g, 193.8 ± 4.80 g and 193.8 ± 4.02 g, respectively.

Toxic effect on food intake

The mean food intake of control and different extracts treated rats during the experimental period was shown in Figure 1. The control rats showed normal food intake throughout the experimental period. The food intake of ethanol, acetone and benzene extract treated rats showed a slight increase in the week I (16.8 ± 1.35 g, 18 ± 2.56 g, 17.7 ± 1.67 g) but it showed decreased levels of food intake in the week II (16.2 ± 19.0 g, 15.5 ± 18.2 g, 16.2 ± 20.5 g) respectively when compared to food intake in week I (19.3 ± 2.59 g), and week II (18.9 ± 2.09 g) of control rats.

Toxic effect on water intake

Mean water intake of control and different extracts treated rats was shown in Figure 1. The control rats showed a normal increase in the water intake throughout the experimental period (22.1 ± 1.85 ml and 22.6 ± 2.57 ml in week I and II respectively). Ethanol, acetone and benzene extracts treated rats showed decreased level of water intake in week II (22.81 ± 1.59 ml, 19.0 ± 23.0 ml and 19.8 ± 23.8 ml, respectively) when compared to the water intake of the rats in initial day (23.7 ± 1.72 ml, 23.5 ± 1.90 ml and 23.9 ± 2.10 ml, respectively). However, the extract treated rats showed the decreased trend of water intake at the end of the experimental period, it showed more or less similar to the water intake of the control rats.

Toxic effect on organ weight

Effect of treatment of the plant extract on relative organ weights are shown in Figure 2. The relative weight of liver of ethanol, acetone and benzene extract treated rats (3.67 ± 0.226 , 3.69 ± 0.229 and 3.71 ± 0.736 g/100g body weight, respectively) were observed to be more or less similar to that of control rats (3.98 ± 0.399 g/100g body weight). The mean relative weight of heart of ethanol, acetone and benzene extract treated rats was observed to be similar to that of control rats (0.4 ± 0.05 g/100g body weight). The mean relative weight of lungs of the ethanol, acetone and benzene extract treated rats showed a slight increase (0.7 ± 0.18 , 0.7 ± 0.28 and 0.7 ± 0.10 g/100g body weight, respectively) when compared to the control rats (0.6 ± 0.09 g/100g body weight). The relative weight of both right and left kidney showed a slight increase when compared to the control rats (0.5 ± 0.06 and 0.5 ± 0.07 g/100g body weight of right and left kidney respectively).

Toxic effect on haematological parameters

Analysis of blood parameters with respect to animal studies has a high relevance and predictive value for human (Rhiouani et al., 2008; Koshy et al., 2011). The assessment of haematological parameters could be used to reveal the deleterious effect of foreign compounds toxins, chemicals and plant extracts on the blood constituents of animals. They are also used to determine possible alterations in the levels of biomolecules such as enzymes, metabolic products, haematology, normal functioning and histomorphology of the organs (Maglhaes et al., 2008; Oyedemi et al., 2011). Haematological parameters of control and extract treated rats are shown in figure 3a and 3b. The total WBC count and the differential count except lymphocytes and basophil of the extract treated rats was decreased when compared to the control rats ($9.02 \pm 0.350 \times 10^3/\mu\text{L}$). Basophil was totally absent in the extract treated rats and the lymphocyte count of benzene extract treated rats was similar ($0.2 \pm 0.005\%$) when compared to control ($0.2 \pm 0.010\%$) and the ethanol, and acetone extract treated rats. The WBC differential count values in the ANOVA table are expressed in arcsine values. The total RBC count, haemoglobin, haematocrit, MCV, MCH, MCHC showed increased levels when compared to control rats but the levels are not exceeded from the normal range. The platelet count of the control and the extract treated rats differed from each other but the values are within the normal range. All the haematological parameters of control, ethanol, acetone and benzene extract treated groups showed significant difference except MCH (one way ANOVA; $p < 0.05$, SNK test).

Acute toxic effect of different extracts of T. chebula on histoarchitecture of vital organs:

Photomicrography of lungs, heart, liver and kidney of control and different extract of plant powder treated rat groups are shown in Plates 1-4. The control and extract treated rats showed normal alveoli, alveolar duct and blood vessels. The normal bronchi lined by ciliated epithelium are observed in the extract treated groups. The muscle of the heart exhibited alternatively light and dark bands and possessed a normal central nucleus in all the extract treated rats. The liver of control rat showed normal hepatic lobules, hepatocytes and central vein. The cell cords were separated by narrow blood sinusoids. Sinusoidal capillaries (sinusoids) separate the sheets of hepatic cells and empty into the central veins. The hepatic cells were thicker and the sinusoids appear as light areas between the cords of cells. The nuclei of hepatic cells were large and spherical, bi-nucleated cells also found. Histological sections of kidney of all the groups showed that the glomeruli, tubules, blood vessels and interstitium appear normal. No pathological changes were observed in test herbal drugs treated rat kidney. There was no macroscopic change of the central organ (such as appearance, colour and size) considered to be related to the treatment. The histopathological examinations of lungs, heart, liver and kidney revealed no remarkable morphological alteration in all treatment and control groups.

IV CONCLUSION

The results of this study showed no changes in the behaviour, no toxic symptoms, and changes in the body weight, food intake, water intake, and relative organ weight. However, the haematological parameters differed from each other but it does not exceed from the normal range. The histoarchitecture of the vital organs did not show any damaged cells, blood vessels and tubules in all the extract treated rats. Thus the present study revealed that the Terminalia chebula fruit powder extract at 2000 mg/kg body weight does not produce any toxic effect in the ethanol, acetone and benzene extract treated rats.

ACKNOWLEDGEMENT

I thank the Management and the Principal, Holy Cross College (Autonomous), Tiruchirappalli for providing necessary facilities, and Department of Science and Technology for financial support.

REFERENCES

- [1]. A. G Agbor, and Y.J. Ngogang, Toxicity of herbal preparations, Cameroon Journal of Ethnobotany, 1, 2005, 23-28.
- [2]. A. J. Akindele, and O. O. Adeyemi, Antiinflammatory activity of the aqueous leaf extract of Byrsocarpus coccineus. Fitoterapia, 78(1), 2007, 25-28.

- [3]. M. A. Ali, J. H. Oh, and P. J. Kim, Evaluation of silicate iron slag amendment on reducing methane emission from flood water rice farming, *Agriculture, ecosystems and environment*, 128(1-2), 2008, 21-26.
- [4]. H. J. Baker, J. R. Lindsey, and S. H. Weisbroth, Housing to control research variables, In: *The Laboratory Rat*, Vol. I.
- [5]. J. Bateman, R. D. Chapman, and D. Simpson, Possible toxicity of herbal remedies. *Scoff. Med. J.*, 1998, 43: 7-15.
- [6]. J. Bruneton, *Plantes Médicinales: Phytochimie, Pharmacognosie*. Lavoisier, New York 914 2, 1993, éme Ed.
- [7]. B.D Doss, G.K Rhoades, S.M Stanley, and H. J. Markman, The effect of the transition to parenthood on relationship quality: an 8-year prospective study. *Journal of personality and social psychology*, 2009, 96 (3): 601.
- [8]. Erhirhie Earnest Oghenesuvwe and Goodies Emuesiri Moke. *Xylopi aethiopica*: A Review of its Ethnomedicinal, chemical and pharmacological properties. *American Journal PharmTech Research*, 2014, 4(6): 21-37.
- [9]. A.Fola, Local medicinal plants and the health of the consumers. A paper delivered at the PSN/CF PCON organization workshop in *Clinical Pharmacy and Herbal Medicine*, 1993, 9: 28-31.
- [10]. J. Harborne, *Phytochemical methods, a guide to modern techniques of plant analysis*, JB Harborne. Chapman. 1973, London. GB.
- [11]. P. Hodek, P. Trefil, and M. Stiborova, Flavonoids - Potent and versatile biologically active compounds interacting with cytochrome P450. *Chemico-Biol. Intern.*, 2002, 139(1): 1-21.
- [12]. P.V. Kadam, R.S. Deod, Shivatare, K.Y. Yadav, and M.J. Patil, Pharmacognostic, phytochemical and physicochemical studies of *Mimusops Elengi* Linn stem bark (Sapotaceae). *Der Pharm Lettr.*, 2012, 4: 607-613.
- [13]. V.P. Kamboj, Herbal medicine. *Journal of Current chemical and pharmaceutical Sciences*, 2000, 78(1): 35-9.
- [14]. Kokate, G. Tushar, B. E Svensson, and M. A. Rogawski, Anticonvulsant activity of neurosteroids: correlation with gamma-aminobutyric acid-evoked chloride current potentiation. *Journal of Pharmacology and Experimental Therapeutics*, 270(3), 1994, 1223-1229.
- [15]. R. K. Koshy, B. Raj Kapoor, and M. Azmathulla, Acute and subacute toxicity of methanol extract of *Elytraria acaulis landau* in rat. *Pharmacology online*, 2011, 3: 229-242.
- [16]. U. Krings, and R. G. Berger, Antioxidant activity of some roasted foods. *Food Chemistry*, 2001, 72: 223-229.
- [17]. P. Magalhaes, H. Appell, and J. Duarte, Involvement of advanced glycation end products in the pathogenesis of diabetes complication: the protective role of regular physical activity. *Evr Rev Aging Phys Act.*, 5(1), 2008, 17-29.
- [18]. C. Mahidol, H. Prawat, V. Prachyawarakorn, and S. Ruchirawat, Investigation of some bioactive Thai medicinal plants. *Phytochemistry Reviews*, 2002, 1: 287-297.
- [19]. D.E. Okwu, and F. N. Morah, Mineral and nutritive value of *Dennettia tripetala* fruits. *Fruits*, 59(6), 2004, 437-442.
- [20]. S. O. Oyedemi, M. T. Yakubu, and A. J. Afolayan, Antidiabetic activities of aqueous leaves extract of *Leonotis leonurus* in streptozotocin induce diabetic rats. *Journal of Medicinal Plant Research*, 5(1), 2011, 119-125.
- [21]. N. Raaman, *Phytochemical techniques*. 2006, New India Publishing.
- [22]. P. Rajesh, S. Latha, P. Selvamani, and V.R. Kannan, Phytochemical screening and toxicity studies on the leaves of *Capparis sepiaria* Linn. *Capparidaceae*. *Journal of basic and clinical pharmacy*, 2010, 1(1).
- [23]. V. Rajpal, *Testing and extraction methods of Medicinal Herbal, Standardisation of Botanicals*. Estern Publishers, 2002, New Delhi 2:124.
- [24]. H. Rhiouani, J. El-Hilaly, Z. H. Israili, and B. Lyoussi, Acute and sub-chronic toxicity of an aqueous extract of the leaves of *Herniaria glabra* in rodents. *Journal of ethnopharmacology*, 118(3), 2008, 378-386.
- [25]. N. Salah, N. J. Miller, G. Paganga, L. Tijburg, G. P. Bolwell, and C. Rice-Evans, Polyphenolic flavanols as scavengers of aqueous phase radicals and as chain-breaking antioxidants. *Archives of biochemistry and biophysics*, 322(2), 1995, 339-346.
- [26]. Singh and Meenakshi, Antimicrobial flavonoid rutin from *Pteris vittata* L. against pathogenic gastrointestinal microflora. *American Fern Journal*, 98.2, 2008, 98-103.
- [27]. A. Sofowora, *Medicinal Plants and Traditional Medicine in Africa*. Spectrum Books Limited, Ibadan, Nigeria, 1993, 151-153.
- [28]. Y.R. Sohini, and R.M. Bhatt, Activity of crude extract formulation in experimental hepatic amoebiasis and in immunomodulation studies. *J Ethnopharmacol.*, 1996, 54: 119-124
- [29]. Ugbogu and Eziuche Amadike. Toxicological assessment of the aqueous dried leaf extracts of *Senna alata* L. in wistar rats. *African Journal of Pharmacy and Pharmacology*, 10(34), 2016, 709-717.

Elavarasi, S." Evaluation of Toxic Effect of Terminalia Chebula Fruit Extract in Albino Rats" International Journal of Pharmaceutical Science Invention(IJPSI), vol. 07, no. 04, 2018, pp. 28-32.