

Identification of the chemical composition of extracts from the trunk bark of *Pseudocedrelakotschyi* and *Khaya senegalensis* by Gas Chromatography-Mass Spectrometry coupling

KoudoroYaya Alain¹, BothonFifaThéomaine Diane^{1,2}, AgbangnanDossaCokou Pascal¹, Alitonou Guy Alain¹, AvlessiFélicien¹ and Sohounhloue C. K Dominique¹

¹Laboratory of Study and Research in Applied Chemistry, Polytechnic School of Abomey-Calavi, University of Abomey-Calavi, 01 PO Box 2009, Cotonou, Benin;

²Kaba Laboratory for Research in Chemistry and Applications, National Higher Institute of Industrial Technology, National University of Sciences Technologies Engineering and Mathematics, Benin

Corresponding Author: KoudoroYaya Alain,

ABSTRACT: The use of plants for therapeutic purposes dates back to the appearance of man in the face of the various adversities of nature. The GC-MS technique was used to characterize metabolites from the stem barks of *Pseudocedrelakotschyi* and *Khaya senegalensis*. Through the results obtained, we note at the level of the bark of *Khaya senegalensis* ten major compounds such as: 4-hydroxyionone, 1,2,4,5-tetramethylbenzene, 11-hydroxyresibufogenine; 17-Hydroxy-3,20-dioxopregna-4,9,11-trien-21-yl acetate; pimara-7,15-dien-3-one; 9,19-cycloergost-24-en-3-ol, 4,14-dimethyl-acetate; cholesterol; lycopene; deacetylcinobufagine; 3-hydroxy-17-oxo-androsta-5,7,9-triene. On the other hand, at the level of the trunk bark of *Pseudocedrelakotschyi*; astaxanthin; campesterol; stigmasterol; sitosterol and stigmast-4-en-3-one have been identified as major compounds. The richness in metabolites of these two plants could explain their use in traditional medicine.

KEYWORDS: *Pseudocedrelakotschyi*, *Khaya senegalensis*, Soxhlet, Extract, Chromatography, Metabolites.

Date of Submission: 01-07-2022

Date of acceptance: 13-09-2022

I. INTRODUCTION

The global market for herbal medicines is continuously expanding, with an annual global value estimated at around US\$800 million (OOAS, 2013). In recent years, human knowledge of medicinal plants has constantly expanded and deepened from one civilization to another (Kamariet *et al.*, 2009). Plants have a great importance on both humans and animals well-being (Bako *et al.*, 2005). They are used in medicine, food, cosmetics and energy. Plants have shown some therapeutic efficacy in the treatment of many pathologies against which modern medicine has remained powerless. *Pseudocedrelakotschyi* (syn: *Cedrelakotschyi*) and *Khaya senegalensis* (*Swietenia senegalensis*), of the family Meliaceae are widely distributed in Africa (Kerharo, 1974). The Meliaceae are large trees with bark that is often fragrant, greyish, dark and scaly and sometimes low in latex (kerharo and Adam, 1974; Shahina and Saranna, 1989). These leaves are alternate, usually pinnately compound, sometimes trifoliate, unifoliate or simple leaves without stipules (Shahina and Saranna, 1989). Their different organs are heavily exploited in traditional medicine. They are used to treat deworming, malaria, fever, jaundice, colic, scabies, leprosy, wound dermatitis, anemia diarrhea and migraine, gastrointestinal diseases, rheumatism, dysentery, epilepsy... (Adjanohoun *et al.*, 1986; Olayinka *et al.*, 1992; Traoré, 1999; Doumbia, 1994; Togola, 2002; Georgewill, 2008; Akande and Hayashi, 1998; Tapsoba and Deschamps, 2006). Previous work has proven that the different organs of *P. kotschyi* and *K. senegalensis* possess antidysenteric, antiulcer, analgesic, anti-inflammatory, antidiabetic, antimalarial, antibacterial and antidiarrheal activities (Asase *et al.*, 2005; Kassimet *et al.*, 2009; Anuka *et al.*, 1999; Musa *et al.*, 2008; Georgewill, 2008; Sablassou, 1996). In the literature, very few works have addressed the characterization of extracts of *Pseudocedrelakotschyi* and *Khaya senegalensis*. In view of the use of these two plants in traditional medicine and their biological activities, it becomes necessary to deepen the characterization of their metabolites.

II. MATERIAL AND METHODS

- Plant collection

Pseudoecdreلاكotschy and *Khaya senegalensis* plants materials were collected from Abomey-calavi and Dassa (Benin)

- Processing of plant samples

The samples of *Pseudoecdreلاكotschy* and *Khaya senegalensis*, once harvested, were dried in the laboratory until their plant mass stabilized before being made into powder

- Extract preparation

One gram of powder from the bark of each plant (*Pseudoecdreلاكotschy* and *Khaya senegalensis*) was introduced into the cartridge of the Soxhlet extractor. 50 mL of dichloromethane were poured into the flask topped with the Soxhlet extractor system. The whole was heated at 40°C for 12 hours and the dichloromethane extract was recovered.

Characterization of the compounds of the trunk barks *Pseudoecdreلاكotschy* and *Khaya senegalensis*:-

The structural determination of molecules of dichloromethane extract of trunk bark of *Pseudoecdreلاكotschy* and *Khaya senegalensis* was made by varian GC/MS type 1200 T, in positive electron impact mode (ionization energy: 70 eV), equipped with a single injector and a VF-5MS type fused silica capillary column (25 m long; 0.25 mm external diameter and 0.25 µm internal diameter). The carrier gas used is helium with a flow rate of 1mL/min which is constant throughout the analysis. The temperature of the injector is 40°C that of the source is 281°C and 298.6°C at the level of the detector and the transfer line. The oven temperature program is defined by an isothermal temperature of 40°C for 5 min with a ramp of 5°C/min up to 310°C, followed by a second isotherm at 310°C for 1min. The quantity of injected extract is 1µL. The injection was made in splitless mode. The mass spectrometry data were established in scan mode whose m/z ratio varies between 50-800 amu. The identification of the compounds present in the extracts was made by comparing the spectra obtained with the information provided by the NIST database (Eswaran *et al.*, 2012; Bojaja *et al.*, 2012; Apostolides *et al.*, 2013).


III. RESULTS AND DISCUSSION

Gas chromatography-mass spectrometry analysis of dichloromethane constituents of trunk barks of *K. senegalensis* and *P. kotschy* led to the identification of several metabolites.

Metabolites from stem bark of *K. senegalensis* and *P. kotschy*

- Metabolites from the trunk bark of *K. senegalensis*

The compounds characterized in the dichloromethane extract of *Khaya senegalensis* are indicated by the chromatogram in figure 1.



FigureNo.1: Chromatographic profile of dichloromethane extract of trunk bark of *K. senegalensis*

In the dichloromethane extract of the trunk bark of *Khaya senegalensis*, 10 major compounds such as 4-hydroxyionone, 1,2,4,5-tetramethylbenzene, 11-hydroxyresibufogenine; 17-Hydroxy-3,20-dioxopregna-4,9,11-trien-21-yl acetate; pimara-7,15-dien-3-one; 9,19-cycloergost-24-en-3-ol, 4,14-dimethyl-acetate; cholesterol; lycopene; deacetylcinobufagine; 3-Hydroxy-17-oxo-androsta-5,7,9-triene have been identified with forty-six minority compounds. The richness and diversity in secondary metabolites of the trunk bark of *K. senegalensis* could justify its use in traditional medicine. An extract of the bark of *Khaya senegalensis* is commonly used in African traditional medicine for pain and inflammation. *Khaya senegalensis* bark extract was hypothesized to contain inhibitors of the cyclooxygenase-2 (COX-2) gene and to be useful in the prevention and treatment of colorectal cancer.

TableNo.1: Metabolites from the trunk bark of *K. senegalensis*

RT (min)	Metabolites
31.524	4-hydroxyionone
39.461	1,2, 4, 5- tetramethylbenzene
55.229	11-hydroxyresibufogenine

55.571	17-Hydroxy-3,20-dioxopregna-4, 9,11-trien-21-yl acetate
55.776	Pimara-7,15-dien-3-one
58.261	9,19-cycloergost-24-en-3-ol, 4,14-dimethyl-acetate
58.641	Cholesterol
58.823	Lycopene
59.178	Deacetylcinobufagine
59.347	3-hydroxy-17-oxo-androsta-5, 7,9-triene
15.076	2,7,7-trimethylbicyclo[2,2,1]hept-2-ene
25.793	Caryophyllene
26.953	4-methyl-1, 2, 3, 4, 4, 5, 6, 7-octahydro-2-naphthalenol
28.129	Cadinene
28.560	1,4-dimethyl-3-tetrahydroacetophenone
29.508	3-methylene-1, 5, 5-trimethylcyclohexene
30.270	1,4-dimethyl-3-tetrahydroacetophenone
32.186	1, 2, 3, 4-tetrahydronaphthalen-1,5-diol
32.397	5-isopropylidene-4,6-dimethylnona-3,6,8-trien-2-ol
33.074	2-octenoic acid, 4-isopropylidene-7-methyl-6-methylene-methyl ester
35.062	Endothaldimethyl ester
37.236	3-methyl-5-phenylhex-1-en-4-ol
37.381	Ambrettolide
37.711	Guaiene
37.846	1, 2, 3,4-tetrahydroisoquinoline
38.848	Tricyclo[4,2,2,0]deca-7-ene
40.310	Sericealactone
41.080	7-methyl-1,4,5,6,7,7hexahydro hind-2-en-2-one
41.182	2, 2-dimethyl-3-vinylbicyclo [2, 2,1] heptane
41.285	Cuminal
41.845	2-[2-methylpropenyl] cyclohexanone
42.513	P-menth-4-en-3-one
43.654	Norfenefrin
45.673	Diphenyl-3,3'-dimethyl-2,2'-ditertiobutyl-1,1'-dihydroxymethane
45.878	Neonicotine
46.081	(Z)-9-tetradecenal
46.749	Methylretinoate
46.842	2-Acetyl-5,8-dihydroxy-3-methoxynaphthoquinone
52.042	retinal
52.397	Ursodiolo
52.786	3-ethyl-3-hydroxyandrostan-17-one
53.623	3-hydroxy-17-oxo-androsta-5,7,9-triene
53.784	Methylretinoate
54.887	Ergosterylacetate
55.057	4,6-cholestadien-3-ol
55.913	3-ethyl-3-hydroxyandrostan-17-one
56.626	2-(7-hydroxymethyl-3,11-dimethyldodeca-2, 6,10-trienyl)-[1,4] benzoquinone
56.737	Resibufogenin
57.194	11-hydroxyresibufogenin
57.583	17-hydroxy-3,20-dioxopregna-1, 4, 9-trien-21-yl acetate
57.761	Carotene
58.057	lupeylacetate
58.184	Sitosterol
58.924	Stigmasta-4, 22-dien-3-one

Legend.RT: Retention Time

• **Metabolites from the trunk bark of *P. kotschy***

Figure 2 shows the chromatographic profile of the dichloromethane extract from the trunk bark of *P. kotschy*.




Figure No.2: Chromatographic profile of dichloromethane extract of stem bark of *P. kotschy*

The compounds identified in dichloromethane extract of trunk bark of *P. kotschyi* are listed in Table 2. Sixteen (16) compounds were identified in dichloromethane extract of trunk bark of *P. kotschyi* with astaxanthin, campesterol, stigmaterol, sitosterol and stigmast-4-yn-3-one the majority compounds. The richness and diversity in secondary metabolites of trunk bark of *P. kotschyi* could justify its use the traditional treatment of various diseases. For example, cholesta-4, 6-dien-3-ol has interesting antimicrobial activity (Yong-Zheet *et al.*, 2018).

Table No.2: Chemical composition of dichloromethane extract from trunk bark of *P. kotschyi*

RT (min)	Metabolites
55.094	Astaxanthin
57.109	Campesterol
57.430	Stigmaterol
58.184	Sitosterol
59.732	Stigmast-4-in-3-one
51.639	3-[3-Methoxy-4-hydroxyphenyl]-2-oxopropanoic acid
53.656	Ursodiol
53.943	Cholesta-4,6-dien-3-ol
54.222	Ergost-5-en-3-ol acetate
54.578	3-ethyl-3-hydroxyandrostane-17-one
55.890	Vitamin E
58.285	Cyclolanost-24-en-3-ol acetate
58.641	Cholest-5-en-3-one
58.945	Cholesta-4,22-dien-3-one
59.114	Ursodeoxycholic acid

Legend. RT: Retention Time

IV. CONCLUSION

The use of herbal medicine is common in Africa to treat various ailments and diseases. This research work has made it possible to identify the metabolites of the bark of *Khaya senegalensis* and *Pseudocedrelakotschyi*, plants that are in high demand in traditional medicine in Benin to treat several diseases. From the results obtained, the trunk barks of *Khaya senegalensis* and *Pseudocedrelakotschyi* are rich and diversified in secondary metabolites. The main metabolites of the bark of *Khaya senegalensis* are 4-hydroxyionone, 1,2,4,5-tetramethylbenzene, 11-hydroxyresibufogenin, 17-hydroxy-3,20-dioxopregna-4, 9,11-trien-21-yl acetate, pimar-7,15-dien-3-one; 9,19-cycloergost-24-en-3-ol, 4,14-dimethyl-acetate; cholesterol, lycopene, deacetylcinobufagine, 3-hydroxy-17-oxo-androsta-5,7,9-triene. At the level of the trunk bark of *Pseudocedrelakotschyi*, astaxanthin, campesterol, stigmaterol, sitosterol and stigmast-4-yn-3-one are the main metabolites. The strong use of *Khaya senegalensis* and *Pseudocedrelakotschyi* in traditional medicine could be explained by their richness and diversification in metabolites.

• Conflicts of Interest

The authors declare that they have no conflicts of interest

REFERENCES:

- [1]. Organisation Ouest Africaine de la Santé (OOAS). Pharmacopée de l'Afrique de l'Ouest, 2020. p.321;
- [2]. Tapsoba H, Deschamps JH. Use of medicinal plants for the treatment of oral diseases in Burkina Faso J of Ethnopharmacol, 2006. 10: P. 68-78
- [3]. Kamari P., Otaghvaria A. M., Govindaparyi H., Bahuguna M., Uniyal P. Some ethno-medically important of India. International Medical Aromatic of Plant, 2009 1, p.18-22;
- [4]. Bako SP, Bakfur MJ, John I, Bala EI. Ethnomedicinal and phytochemical profile of some savanna plant species in Nigeria. Int J Bot, 2005; p. 147-150;
- [5]. Kerharo J., Adam J.G. (1974). La Pharmacopée sénégalaise traditionnelle: Plantes médicinales et toxiques. Edition Vigot frères. Paris, 3-545.
- [6]. Adjanohoun E, Adjakidjè V, Ahyi MRA, Aké Assi L, Akoègninou A, D'Almeida J, Apovo F, Boukef K, Chadaré F, Cusset G, Dramane K, Eyme J, Gassita J-N, Gbaguidi N, Goudoté E, Guinko S, Hounnon P, Issa L, Keita A, Kiniffo HV, Koné Bamba D, Musampa Nseyya A, Saadou N, Sodogandji T, DE Souza S, Tchabi A, Zinsou Dossa C, Zohoun T. 1989. Contribution aux Etudes Ethnobotaniques et Floristiques en République Populaire du Bénin. ACCT: Paris.
- [7]. Olayinka, A.I. and Olorunfemi, M.O. Determination of Geoelectrical Characteristic in Okene Area and Implication for Boreholes. Journal of Mining and Geology, 1992.28: P. 403-412;
- [8]. Doumbia L Les effets de *Melia Azedarach* sur les larves du criquet pèlerine *Schistocerca gregaria* Forsk. Sahel Protection des végétaux. INFO, 1994. 60: 2-10.
- [9]. Akande John A, Hayashi and Yamamoto Koichi. Extractive components of tropical chewing stick species. Trop. Sci., 1998, 38: p.87-90;

Identification of the chemical composition of extracts from the trunk bark of ..

- [10]. Asase Alex, Alfred A Oteng-Yeboah, George T Odamttten, Monique S J Simmonds. Ethnobotanical study of some Ghanaian anti-malarial plants. *J Ethnopharmacol.* 2005 99(2):273-279.
- [11]. Togola A (2002). Etude de la phytochimie et de l'activité antipaludique de *Alchorneacordifolia* Schumacher. These de pharmacie Bamako, 76P;
- [12]. Georgewill OA and Georgewill UO (2008). Antiarthritic activity of *pseudocedrelakotschy* in albino rats. *African Journal of Applied Zoology & Environmental Biology*, 10:70-72;
- [13]. Tapsoba H, Deschamps JH. Use of medicinal plants for the treatment of oral diseases in Burkina Faso *J of Ethnopharmacol*, 2006. 10: P.68-78.
- [14]. Eswaran R, Anandan A, Doss A, Sangeetha G and Anand SP. Analysis of chemical composition of *cissusquadrangularis* Linn. By GC-MS. *Asian Journal of Pharmaceutical and Clinical Research*, 2012. 5(2): P.139-140.
- [15]. Bojaxa AR and Rosakutty PJ (2012). GC-MS analysis of methanol wild plant and callus extracts from three *Cissus* species, Family Vitaceae. *Journal of Chemical and Pharmaceutical Research*, 4(7): 3420-3426;
- [16]. Apostolidis NA, Beyrouthy MEI, Dhifi W, Najm S, Cazier F, Najem W, Labaki M and AbouKaïs A. Chemical Composition of Aerial Parts of *Rosmarinus officinalis* L. Essential Oil Growing Wild in Lebanon. *Journal of Essential Oil Bearing Plants*. 2013. 16 (2): P. 274-282
- [17]. Yong-Zhe Zhu, Jing-Wen Liu, Xue Wang, In-Hong Jeong, Young-Joon Ahn and Chuan-Jie Zhang ORCID (2018). Anti-BACE1 and Antimicrobial Activities of Steroidal Compounds Isolated from Marine *Urechisunicinctus*, 16(3), pp: 1-12
- [18]. Musa YM, Haruna AK, Ilyas M, Yaro AH, Ahmadu AA, Usman H. Phytochemical, analgesic and anti-inflammatory effects of the ethyl acetate extract of the leaves of *Pseudocedrelakotschy*. *Afr. J. Trad. Compliment. Alt. Med*, 2008. 5:92-96.
- [19]. Anuka, J.A. Ijezie, D.O., & Ezebnik, O.N. "Investigation of Pharmacological actions of the extract *Pseudocedrelakotschy* in Laboratory animals. ABSTRACTS of the proceedings of XXVIIth Annual Regional Conference of WASP, 1999: P. 9-10.
- [20]. Togola A. Etude de la phytochimie et de l'activité antipaludique de *Alchorneacordifolia* Schumacher. These de pharmacie Bamako, 2002. 76P,
- [21]. Rani Thakur, R. And A. Verma, Mouth Dissolving Tablets- Preparation Characterization And Evaluation: An Overview. *Journal Of Pharmacy Research*, 2012. 5(2): P. 993-1000.
- [22]. Sreenivas, S., Et Al., Orodispersible Tablets: New-Fangled Drug Delivery System-A Review. *Indian Journal Of Pharmaceutical Education*, 2005. 39(4): P. 177.
- [23]. Ghosh, T. A. Ghosh, And D. Prasad, A Review On New Generation Orodispersible Tablets And Its Future Prospective. *International Journal Of Pharmacy And Pharmaceutical Sciences*, 2011. 3(1): P. 1.
- [24]. Chue, P. R. Welch, And C. Binder, Acceptability And Disintegration Rates Of Orally Disintegrating Risperidone Tablets In Patients With Schizophrenia Or Schizoaffective Disorder. *Canadian Journal Of Psychiatry*, 2004. 49(10): P. 701-703.
- [25]. Seager, H. Drug Delivery Products And The Zydys Fast Dissolving Dosage Form*. *Journal Of Pharmacy And Pharmacology*, 1998. 50(4): P. 375-382.
- [26]. Dixit, S. Et Al. Fast Dissolving Tablet-A Promising Approach For Drug Delivery: A Review. *Journal Of Pharmacy Research*, 2012. 5(3): P. 1508-1513.
- [27]. L Lachman, HA Liberman, Joseph L Kani G, The theory and practice of industrial pharmacy, Varghese publishing house, Bombay, 1990, 3rd Edition, 315-317.
- [28]. Kuchekar, B. A.C. Badhan, And H. Mahajan, Mouth Dissolving Tablets: A Novel Drug Delivery System. *Pharma Times*, 2003. 35(7).
- [29]. Harmon, T. Emerging Technology Beyond The First Generation Of Orally Disintegrating Tablets September 1, 2006.
- [30]. Chang, R.-K., Et Al., Fast-Dissolving Tablets. *Pharmaceutical Technology*, 2000. 24(6): P. 52-58.
- [31]. Siddiqui, M.N. G. Garg, And P.K. Sharma, Fast Dissolving Tablets: Preparation, Characterization And Evaluation: An Overview. *International Journal Of Pharmaceutical Sciences Review And Research*, 2010. 4(2): P. 87-96.
- [32]. Fu, Y. Et Al. Orally Fast Disintegrating Tablets: Developments, Technologies, Tastemasking And Clinical Studies. *Critical Reviews™ In Therapeutic Drug Carrier Systems*, 2004. 21(6).
- [33]. T Shu, H Suzuki, K Hironaka, K Ito, *Chem Pharm Bull.*, 2002, 50(2), 193-198.
- [34]. J Madan, AK Sharma, R Singh, *Trop J Pharm Res.*, 2009, 8(1), 63-70.