

## Contribution to the phytochemical and biological study of the methanolic extract of root bark of *Mezoneuron benthamianum* Baill. : a medicinal plant from Ivory Coast

Ouattara Logopho Hyacinthe<sup>1,2\*</sup>, N'guessan Patrick Audrey<sup>2</sup>, Ambeu-Loko N'ta Christelle Mélissa<sup>3\*</sup>, Sima Obiang Cédric<sup>4</sup>, Philippe Jéhan<sup>5</sup>, Fabian Lambert<sup>5</sup>, Nicolas Le Yondre<sup>5</sup>, Kablan Ahmont Landry Claude<sup>1</sup>, Mamyrbékova-Békro Janat Akhanovna<sup>2</sup>, Békro Yves-Alain<sup>2</sup>

<sup>1</sup>UPR of organic chemistry, Department of Mathematics-Physics-Chemistry, UFR Biological sciences, Peleforo Gon Coulibaly University, Korhogo, Ivory Coast

<sup>2</sup>Laboratory of Bio-Organic Chemistry and Natural Substances, UFR Fundamental and Applied Sciences, Nangui Abrogoua University, Abidjan, Ivory Coast

<sup>3</sup>Laboratory of Constitution and Reaction of Matter, UFR Structural Sciences of Matter and Technology, Félix Houphouët-Boigny University, Abidjan, Ivory Coast

<sup>4</sup>Biochemistry Research Laboratory (LAREBIO), Faculty of Sciences, Masuku University of Science and Technology, Franceville, Gabon

<sup>5</sup>Western Regional Center for Physical Measurements-CRMPO, ScanMAT UAR 2025 CNRS, University of Rennes 1, Beaulieu Campus, 263 Avenue du General Leclerc, CS 74205, 35042 Rennes Cedex, France

Corresponding Authors: Ouattara Logopho Hyacinthe, olhy@live.fr / Ambeu-Loko N'ta Christelle Mélissa, chris\_meliss@yahoo.fr

---

**ABSTRACT:** The identification of molecules in the methanolic extract of *Mezoneuron benthamianum* root bark harvested in the middle of Ivory Coast was carried out by LC-MS. The corresponding results revealed the identification of six (06) phytochemicals, among which the pharmacological properties of five were highlighted. These are 2,4,6-Triallyloxy-1,3,5-triazine, 1,6-bis-O-galloyl-beta-D-glucose, epigallocatechin gallate, 1,4,6-tri-O-galloyl-beta-D-glucose, and resveratrol. Biological tests carried out on this extract have shown that it manifests not only significant anticoagulant and analgesic activities, but also relatively low toxicity. Therefore, the biological potential of methanolic extract from the root bark of *M. benthamianum* could be related to its phytochemical profile.

**KEYWORDS:** *Mezoneuron benthamianum*, analgesic, anticoagulant, toxicity and LC-MS.

---

Date of Submission: 03-08-2024

Date of acceptance: 15-09-2024

---

### I. INTRODUCTION

*Mezoneuron benthamianum* is a plant in the Caesalpiniaceae family. It is a climbing and thorny shrub that is widespread from Senegal to Nigeria. This plant is also present in Ivory Coast where it is found in all plant formations but preferably in secondary formations. Known in Nigeria as "Jenifinran" [1], in Ivory Coast, *M. benthamianum* is called "Ako bohuie" in Krobou; "Akpaba wun" in Akyé; "Ako owué" in Baoulé and "Còlè" in Djimini. This plant is commonly used in folk medicine. In West Africa, it is used to treat malaise, wounds, urethral discharge, ulcers, skin infections, pain, fever, diarrhea, persistent coughs and headaches [1,2,3,4]. In Ivory Coast, the fresh root of *M. benthamianum* is consumed in the form of a toothpick against sexual impotence [5]. Several studies have highlighted the pharmacological activities of the plant. The vine is used as an antiseptic, detergent and healer of torpid wounds [6]. Extracts of root bark have been tested for its vasothoven, antioxidant, aphrodisiac and antibacterial properties [7,8]. Leaf extract has also shown antimicrobial effects [9]. Thus, this study is part of a development strategy contributing to the valorization of medicinal plants in Ivory Coast. Thus, it will specifically evaluate the analgesic, anticoagulant and toxicological activities of the methanolic extract of the root bark of *M. benthamianum* on the one hand and contribute to the determination of the chemical composition of that extract on the other hand, using liquid chromatography coupled with mass spectrometry (LC-MS).

## II. MATERIAL AND METHODS

### MATERIAL

#### Plant material

The root bark of *Mezoneuron benthamianum* was collected in July in Dabakala (8° 21' 48" North, 4° 25' 43" West), a town located in the middle of Ivory Coast. After authentication at the herbarium of the National Floristic Center (CNF) in Abidjan (5° 20' 11" North, 4° 01' 36" West), the bark was cleaned and dried first of all away from the sun for 2 days. Then, they were dried under air conditioning (18°C) for 7 days and in an oven (45°C) for 3 days. The dry bark was then ground into powder using an electric grinder (RETSCH, type SM 100). The resulting vegetable powder was used to prepare the extract to be analyzed.

#### Technical material, chemical and biological products

Technical material was composed of an electric shredder (RETSCH, type SM 100), a permanent stirrer, an oven, a rotary evaporator (BÜCHI Waterbath B-480), a centrifuge, a spectrophotometer, a Thermo Scientific Orbitrap Exploris 480 mass spectrometer, EDTA tubes, a feeding probe, syringes and standard laboratory glassware. As for the chemical and biological products, they consist of methanol (80%), water, acetonitrile (CH<sub>3</sub>CN), acetic acid (1%), physiological water, paracetamol, escin, sodium chloride and sheep's blood.

#### Animal material

For the biological tests *in vivo*, female mice of the species *Mus musculus* weighing between 22 and 32 g, albino of the Swiss breed and from IPCI (Institute Pasteur of Ivory Coast) were used.

### METHODS

#### Extraction

A mass of 50 g of powder was macerated in 375 mL of methanol (80%) for 24 h, at room temperature under permanent stirring. After vacuum filtration, the maceration was concentrated with a rotary evaporator (BÜCHI Waterbath B-480). Then, the concentrate was dried in an oven at 50°C for 48 hours. The crude methanolic extract obtained was stored in the refrigerator at 4°C in a hermetically sealed jar. It was subsequently used to perform analgesic, anticoagulant, toxicity and spectral analysis by LC-MS.

#### Analgesic test

The analgesic test of the methanol extract of *M. benthamianum* root bark was carried out according to the methods described by Koster *and al.* [10] and Chatter *and al.* [11]. Three (3) batches of four (4) mice were made. The blank control batch received 0.1 mL of physiological water. The mice from the other two groups received respectively 0.1 mL of plant extract and 0.1 mL of paracetamol (reference analgesic) at 200 mg/kg BW. Half an hour after administration of the products (vegetable, paracetamol, physiological water), the animals received 0.1 mL of 1% acetic acid (CH<sub>3</sub>COOH) by intraperitoneal injection. The number of abdominal contortions was counted in each mouse for 30 min. The analgesic effect of the plant extract and paracetamol was evaluated by determining the percentage of inhibition of contortions according to the following formula:

$$\% \text{ inhibition} = \frac{\text{Mtb} - \text{Mext}}{\text{Mtb}} \times 100$$

**Mtb**: Average number of contortions of mice from the white control group

**Mext**: Average number of contortions of mice treated with plant extract and paracetamol

#### Anticoagulant test

The study of anticoagulant activity was carried out following the method described by De Freitas *and al.* [12]. Different concentrations (5, 4, 3, 2 and 1 mg/mL) of the methanol extract of *M. benthamianum* and escin (triterpene saponin), were prepared in 0.85% NaCl physiological solution (m/v). The pure NaCl solution constituted the negative control while escin was used as a positive control. These different solutions were subsequently mixed in EDTA tubes with 25 µL of fresh blood from sheep fasted overnight. The mixture was homogenized, incubated for 30 min at 37°C and centrifuged at 3500 rpm for 10 min. During centrifugation, erythrocyte lysis may or may not be observed. The absorbance of the supernatants was measured at 540 nm. The absorbance values as a function of the NaCl concentration enabled to draw a sigmoidal regression curve given by Boltzmann formula, by means of which the percentage of hemolysis of each extract was calculated according to the following formula:

$$\text{Hemolysis (\%)} = \frac{A1 - A2}{1 + e^{\frac{(s-H50)}{ds}}} + A2$$

**A1:** Maximum average absorption values of the sigmoid curve;  
**A2:** Minimum average absorption values of the sigmoid curve;  
**S:** NaCl concentration;  
**H50:** NaCl concentration that causes 50% of hemolysis;  
**ds:** amplitude of the sigmoid curve of the transition between A1 and A2

### Toxicity test

The toxicity study was realized in accordance with OECD Test Guideline 423 for chemical testing [13]. Mice in batches of three (3) were acclimated to the experimental environment in clean, well-ventilated cages for one week before experimentation. During this period, the animals were monitored daily to assess their mass and behaviour. Four (4) batches of these mice, previously fasted for 16 h, were administered 1 mL of *Mezoneuron benthamianum* plant extract by gavage, by administration of variable doses ranging from 300 to 2000 mg/kg bw. After administration of the extracts, the animals were observed for 14 days. The 50% lethal dose (LD50) was determined from the mortality rate of mice.

### LC-MS analysis

The sample was analyzed in nanoLCMS (nLCMS) mode on a Thermo Scientific Orbitrap Exploris 480 spectrometer in positive and negative Electrospray modes. The crude methanolic extract of *Mezoneuron benthamianum* was solubilized in the HPLC grade H<sub>2</sub>O/CH<sub>3</sub>CN (95/5) solvent mixture. The analyses were performed on a quantity of 300 ng of material deposited on the PepMap Neo C18 5µm 300µm x 5 mm / EASY - Spray column PepMap Neo 1500 bars, 75 µm x 150 mm, C18, 2 µm, 100 Å. Positive and negative mode analyses were performed over a 40-minute run with the following instrumental and gradient parameters: CRMPO-PJ-P-480k-40min-100-1500-2000V-SL75-Gradient 2 - Trap and Elute – BackFlush. The interpretations for the positive and negative modes were carried out on a technical triplicate.




Figure 1 : Gradient used for analyzes in positive and negative Electrospray modes


### Statistical analyzes

The analyzes of the measurements obtained during the anticoagulant test were carried out using the GRAPH PAD PRISM software. This software made it possible to determine H<sub>50</sub> and ds parameters.

## III. RESULTS AND DISCUSSION

### Analgesic test


Figure 1 highlights the effect of the methanolic extract of *Mezoneuron benthamianum* root bark on abdominal contractions. The pain inhibition percentage of *M. benthamianum* extract is 47.92% while that of paracetamol is 100%.



**Figure 2 :** Analgesic effect of methanolic extract of *M. benthamianum* roots on abdominal contortions

### Anticoagulant test

Figure 2 presents the anticoagulant effect of the methanolic extract of *M. benthamianum* root bark and the negative and positive controls. The hemolysis percentages of the root bark of *M. benthamianum* vary between  $10.88 \pm 0.26$  and  $17.19 \pm 0.43\%$  while those of escin oscillated between  $21.6 \pm 0.84$  and  $41.13 \pm 0.97\%$ . As for the negative control (physiological NaCl solution), it indicated a hemolysis percentage of 4.47%.



**Figure 3:** Anticoagulant effect of *M. benthamianum* extract, escin and NaCl at different concentrations

### Toxicity test

The results of the toxicity test are recorded in Table 1. This table shows the mortality rate of mice at the different doses of *M. benthamianum* methanolic extract injected. The mortality rate was then used to determine the lethal dose (LD50) of the methanolic extract of the root bark of the plant studied, which is between 500 and 2000 mg/kg bw.


**Table 1 :** Mortality rates of mice as a function of administered doses of *M. benthamianum* methanol extract

Batch mice		Batch 1	Batch 2	Batch 3	Batch 4
Dose administered (mg/kg PC)		300	300	2000	2000
Number deaths per day	1 <sup>st</sup> day	0	0	0	1
	2 <sup>nd</sup> day	1	0	0	0
	3 <sup>rd</sup> day	0	0	0	0
	4 <sup>th</sup> day	0	0	0	0
	5 <sup>th</sup> day	0	0	0	0
	6 <sup>th</sup> day	0	0	0	0


	7 <sup>th</sup> day	0	0	0	0
	8 <sup>th</sup> day	0	0	0	1
	9 <sup>th</sup> day	0	0	0	0
	10 <sup>th</sup> day	0	0	0	0
	11 <sup>th</sup> day	0	0	0	0
	12 <sup>th</sup> day	0	0	0	0
	13 <sup>th</sup> day	0	0	0	0
	14 <sup>th</sup> day	0	0	1	0
<b>Mortality</b>		1	0	1	2
<b>Mortality rate (%)</b>		33.33	0	33.33	66.66

### LC-MS analysis

The chromatographic profiles of *Mezoneuron benthamianum* extract in positive and negative electrospray modes are shown in Figures 4 and 5, respectively. A large number of compounds were highlighted in each of the two chromatographic profiles of the extract.



RT: 0.00 - 40.00


**Figure 5** : Chromatographic profile of *M. benthamianum* extract in negative mode

The formulas of several ions contained in the sample have been highlighted (see supplementary data). The choice was made for the most intense ions, i.e. the most easily ionizable. The LCMS runs were performed at the highest experimental resolution of the Exploris 480 (480000 at mass  $m/z$  200) in order to remove any ambiguity relative to the validation of the raw formula of each ion concerned. In addition, the proposed formulas were validated by modeling theoretical isotopic massifs at experimental resolution considering a maximum experimental error of 2 ppm. From these initial data, a second series of LCMSMS recordings was made in DDA (Data Dependent Acquisition) mode in both polarities in order to access as far as possible the fragmentation spectra of a large number of species. In order to validate the structure of potentially present compounds, the exploitation of these experiments was carried out by comparing the experimental MSMS spectra with those contained in two databases, MassBank (<https://massbank.eu/MassBank/>) and mzCLOUD (<https://www.mzcloud.org/>). Thus, six (6) compounds were identified in positive and negative electrospray modes (Tables 2, 3 and Figure 6). Information on these compounds is reported in Tables 2 and 3, and their molecular structures are shown in Figure 6.


**Table 2**: Possible compounds in *M. benthamianum* extract in positive mode

Retention time (min)	Exact mass of ion [M+H] <sup>+</sup>	Formula of compound	Main fragments of the [M+H] <sup>+</sup> ion observed in MSMS	Database m/zCLOUD	Possible compounds
13.35	479.0820	C <sub>21</sub> H <sub>18</sub> O <sub>13</sub>	-	461.0716, 309.0605, 153.0181	3,4-Di-O-galloylshikimic acid
33.44	250.1186	C <sub>12</sub> H <sub>15</sub> N <sub>3</sub> O <sub>3</sub>	81.0699, 79.0542, 69.9923, 56.0131	-	2,4,6-Tris(allyloxy)-1,3,5-triazine


**Table 3**: Possible compounds in *M. benthamianum* extract in negative mode

Retention time (min)	Exact mass of ion [M-H] <sup>-</sup>	Formula of compound	Main fragments of the ion [M-H] <sup>-</sup> observed in MSMS	Database m/zCLOUD	Possible Compounds
11.07	484.0859	C <sub>20</sub> H <sub>20</sub> O <sub>14</sub>	423.0569, 331.0671, 313.0565, 271.0461, 241.0356, 211.0249, 193.0143, 169.0143, 125.0244	331.0666, 313.0561, 271.0456, 241.0349, 211.0245, 169.0139, 125.0241	1,6-bis-O-galloyl-beta-D-glucose
13.72	457.0776	C <sub>22</sub> H <sub>18</sub> O <sub>11</sub>	305.0667, 169.0142, 125.0243	-	Epigallocatechine


					gallate
13.72	635.0890	C <sub>27</sub> H <sub>24</sub> O <sub>18</sub>	591.0986, 483.0760, 465.0671, 423.0566, 313.0566, 169.0143	591.0986, 483.0774, 465.0668, 439.0877, 423.0565, 331.0667, 313.0561, 295.0456, 169.0139	1,4,6-tri-O-galloyl-beta-D-glucose
21.08	227.0714	C <sub>14</sub> H <sub>12</sub> O <sub>3</sub>	185.0607, 143.0501	-	Resveratrol




3,4-Di-O-galloylshikimic acid




2,4,6-Tris(allyloxy)-1,3,5-triazine




1,6-bis-O-galloyl-beta-D-glucose



Epigallocatechin gallate



1,4,6-tri-O-galloyl-beta-D-glucose



Resveratrol

**Figure 6:** compounds identified in *M. benthamianum* extract

#### IV. DISCUSSION

The present study consisted of analgesic, anticoagulant, toxicity tests and LC-MS spectral analysis of the methanolic extract of *M. benthamianum* root bark. The analgesic test showed that the extract significantly inhibited pain ( $47.92 \pm 0.34\%$ ) compared to paracetamol ( $100 \pm 0.00\%$ ). This pharmacological suitability of *M. benthamianum* root bark could be justified by its phytochemical composition. This organ contains sterols, terpenes, saponins, flavonoids, coumarins, tannins, phenolic acids and alkaloids [3,17]. Several studies have

shown the analgesic effects of some of these secondary metabolites. This is the case for terpenes [18], flavonoids [19], saponins [20], alkaloids [21] and coumarins [22]. These bioactive phytochemicals inhibit the release of mediators (serotonin, acetylcholine, histamine, bradykinin, substance P and prostaglandins (PGE<sub>2α</sub>, PGF<sub>2α</sub>)) involved in the pain process. Similarly, the anticoagulant activity of that extract observed at a haemolysis percentage of 17.19±0.43% is also due to the co-presence of those secondary metabolites. Indeed, bioactive phytochemicals, such as saponins and coumarins, have anticoagulant properties that are widely described in several studies [20, 23, 24]. The manifestation of the anticoagulant effect of the methanol extract of *M. benthamianum* root bark could explain its use in the traditional treatment of hemorrhoidal pathology [17]. Concerning the toxicity test, the treated mice showed some clinical signs such as increased heart rate, breathing difficulties and seizures. However, the animals that survived returned to a normal appearance in the following days. In addition, the death of a few mice was observed after administration of doses between 300 and 2000 mg/kg bw. As a result, according to the OECD guideline [13], *M. benthamianum* root bark would be classified as Category 4, with LD<sub>50</sub> ranging from 500 to 2000 mg/kg bw. Thus, according to the Hodge & Sterner toxicity scale [25], *M. benthamianum* root bark had low toxicity, thus the justification for its recurrent use in traditional medicinal practice. Positive and negative analysis of the methanolic extract of *M. benthamianum* root bark resulted in the structural identification of six compounds. This result subsequently led to the search for their potential biological activities. A systematic review of the literature to support the biological activity of 3,4-Di-O-galloylshikimic acid did not lead to conclusive results. However, the other five molecules identified contain many pharmacological properties. These are 2,4,6-trialloxy-1,3,5-triazine, resveratrol, epigallocatechin gallate, 1,4,6-tri-O-galloyl-beta-D-glucose and 1,6-bis-O-galloyl-beta-D-glucose ; most of them are polyphenols and thus of great therapeutic interest currently:

- 2,4,6-Triallyloxy-1,3,5-triazine is a natural product with antimicrobial activity whose bioactivity has been validated by molecular docking of four antibacterial proteins [26].
- Resveratrol is a polyphenol of the stilbene class found in some plants and also in some fruits [27]. It is a substance induced by environmental or pathogenic stress and intended to contain the damage of the pathogen locally [28]. It has an antiproliferative effect on cultured cancer cells [29] and decreases inflammation in macular degeneration, an age-related vision pathology [30]. Resveratrol is also known as an insulin-sensitizing agent and resistance to weight gain, which also mimics the benefits of physical activity on the remodeling of muscle fibers while increasing resistance to exercise [31].
- Epigallocatechin gallate is a flavanol formed by esterification of epigallocatechin and gallic acid. It is the most abundant flavanol in tea and is known to be a powerful antioxidant [32,33]. It plays a chemopreventive role in ovarian cancer through the regulation of molecular pathways that induce the stem phenotype of ovarian cancer stem cells [34].
- 1,4,6-tri-O-galloyl-beta-D-glucose and 1,6-bis-O-galloyl-beta-D-glucose are two compounds called galloylglucoses. Galloylglucoses that belong to the phenolic acid derivatives and the subclass of gallotannins, are composed of halves of glucose and gallic acid by esterification or O-acylation, and can be divided into mono-, di-, tri-, tetra-, penta-galloylglucose and so on, depending on the number of gallic acid groups. They are considered as a natural defense against herbivores. Over the past few decades, studies have revealed the many biological properties of these compounds. They contain antioxidant, antitumor, anti-inflammatory, antibacterial properties and cardiovascular protective effects, among other properties [35]. Thus, the presence of these different organic compounds in the root bark of *M. benthamianum* could also justify its common use in traditional medicine.

## V. CONCLUSION

In order to enhance the value of the Ivorian flora in general and in particular to provide a rational response to the traditional use of *Mezoneuron benthamianum*, biological investigations (analgesic, anticoagulant and toxicity activity) and chemical investigations (LC-MS analyses) were carried out on the root bark of the said plant. Biological tests showed that the root bark of *M. benthamianum* showed a significant analgesic effect (47.92%) compared to paracetamol (100±0.00%). The non-negligible hemolytic power of this organ was noted with levels between 10.88±0.26 and 17.19±0.43%. The lethal dose of between 500 and 2000 mg/kg bw proves that the root bark of *M. benthamianum* is of low toxicity and therefore does not present any major danger to the body. The phytochemical analysis by LC-MS of the methanol extract of the root bark of *M. benthamianum* highlighted the presence of six phytochemicals, among which pharmacological properties of five were highlighted. These are 2,4,6-Triallyloxy-1,3,5-triazine, resveratrol, epigallocatechin gallate, 1,4,6-tri-O-galloyl-beta-D-glucose, and 1,6-bis-O-galloyl-beta-D-glucose. On the one hand, the biological and phytochemical analyses carried out on *M. benthamianum* constitute a contribution to the valorization of the Ivorian species and on the other hand, justify its use in traditional medicinal practice against some pathologies.

## REFERENCES



- [1]. Irvine R. Woody Plants of Ghana. 1961. Oxford University Press, London : P. 439–481.
- [2]. Dickson R.A., Houghton P.J., Hylands P.J., Gibbons S. Antimicrobial, resistance-modifying effects, antioxidant and free radical scavenging activities of *Mezoneum benthamianum* Baill. *Securinea virosa* Roxb. & Wlld and *Microglossa pyrifolia* Lam. 2006. *Phytotherapy Research*, 20 : P. 41–45.
- [3]. Osho, A. Ethnopharmacological Properties of *Caesalpinia benthamiana*-A Mini Review. 2013. *British Microbiology Research Journal*, 4(2), P. 206-213.
- [4]. N'guessan K. Contribution to the ethnobotanical study in Krobou country, 3rd cycle Doctoral thesis. 1995. National University of Ivory Coast: P. 557.
- [5]. Békro Y. A., Békro M. J. A., Boua B. B., Tra bi F. H., Ehile E. E. 2007. Ethnobotanical study and phytochemical screening of *Caesalpinia benthamiana* (Baill.) Herend. and zarucchi (Caesalpiniaaceae). *Sciences and Nature*, 4 : P. 217-225.
- [6]. Kerharo J., Adam J.G. Traditional Senegalese pharmacopoeia, medicinal and toxic plants. 1974. *Journal of Agriculture and Applied Botany*, Vigot and brothers, Paris (France), P. 548-1011.
- [7]. Dickson, R. A., Houghton, P. J., & Hylands, P. J. Antibacterial and antioxidant cassane diterpenoids from *Caesalpinia benthamiana*. 2007. *Phytochemistry*, 68(10), P. 1436-1441.
- [8]. Zambélé A., Martin-nizard F., Sahpaz S., hennebelle T., Staels B., Bordet R., Duriez P., Brunet C., Bailleul F. Vasoactivity, antioxidant and aphrodisiac properties of *Caesalpinia benthamiana* roots. 2008. *Journal Ethnopharmacology*, 116 (1) : P. 112.
- [9]. Cordell G.A., Binutu O.A. Gallic acid derivatives from *Mezoneuron benthamianum* leaves. 2000. *The Pharmacist Biologist*, 38 : P. 284–286.
- [10]. Koster R., Anderson M., Beer E.J. Acetic acid for analgesic screening. *Federation*. 1959. *Proceedings*. 18 : P. 412-417.
- [11]. Riahi, R. C., Tarhouni, S. S., & Kharrat, R. R. Screening of the anti-inflammatory and analgesic effect of marine algae from the Mediterranean Sea. 2012. *Archives of Pasteur Institute of Tunis*, 88 (1-4), P. 19-28.
- [12]. De Freitas M.V., Netto R.C.M., Huss J.C.C., Souza T.M.T., Costa J.O., Firmino C.B., Penha-Silva N. Influence of aqueous crude extracts of medicinal plants on the osmotic stability of human erythrocytes. 2008. *Toxicology in Vitro*, 22 (1) : P. 219-224.
- [13]. OCDE. Guidelines for the testing of chemical products. 2001. P. 14.
- [14]. Tong H., Bell D., Tabei K., Siegel M. M. Automated Data Massaging, Interpretation and E-Mailing Modules for High Throughput Open Access Mass Spectrometry. 1999. *J. Am. Soc. Mass Spectrom.* 10 : P. 1174.
- [15]. Guo X., Bruins A. P., Covey T. R. Characterization of typical chemical background interferences in atmospheric pressure ionization liquid chromatography-mass spectrometry. 2006. *Rapid Commun. Mass Spectrom.* 20 : P. 3145–3150.
- [16]. Keller B. O., Sui J., Young A. B., Whittall R. M. Interferences and contaminants encountered in modern mass spectrometry. 2008. *Analytica Chimica Acta*. 627 : P. 71-81.
- [17]. Ouattara L.H., Kabran M.G.R., Kadja A.B., Tano B.M., Mamyrbekova-Bekro J.A., Bekro Y.A. Phytochemical study and antioxidant activity of plant extracts from Ivory Coast used in the traditional treatment of hemorrhoids. 2016. *Internationnal Journal of Innovation and Applied Studies*, 15(4) : P. 881-893.
- [18]. Bruneton J., Pharmacognosie, Phytochimie, Plantes médicinales. 1999. 3<sup>ème</sup> Edition, Lavoisier, International EM, Tec. & Doc, Paris, France, P. 1120.
- [19]. Bent H., Havsteen. The biochemistry and medical significance of the flavonoids. 2002. *Pharmacology & Therapeutics*, 96: P. 67–202.
- [20]. Amzal H. Study of the antioxidant activity of saponins in argan cake. 2010. Rabat : Université Mohammed V Agdal. Faculté des Sciences, Rabat ; P. 149.
- [21]. Zenk M., Jueng M. Evolution and current status of the phytochemistry of nitrogenous compounds. *Phytochemistry*, 2007. 68: P. 2757-2772.
- [22]. Reddy N.S., Gumireddy K., Mallireddigari M.R. Novel coumarin-3-(N-aryl)carboxamides arrest breast cancer cell growth by inhibiting ErbB-2 and ERK1. 2005. *Bioorganic and Medicinal Chemistry Letters*, 13 : P. 3141-3147.
- [23]. Ladyguina E., Safronitch L., Otriachenkova V., Bolandina I., Grinkevitch N., Sorokina A., Glizin V.I., Molodjnikova L.M., Mitin Y.S., Samilina I.A. & Ermakova V.A. Analyse chimique des plantes médicinales. 1983. Edition Moska; Vischaya Chkola (traduit du russe): P. 46-347.
- [24]. Park J.D., Rhee D.K., Lee Y.H. Biological activities and chemistry of saponins from *Panax ginseng* C. A. 2005. *Meyer Phytochemistry Reviews*, 4 : P. 159–175.
- [25]. Hodge A.C., Sterner J.H., Toxicity studies: some basic data (A.K DONE). 1980. *Tempo Medical Afrique*, 7 : P. 18.
- [26]. Jeyanthi G., Gnana S. C., Shiny C. L., Deephin T. J. D., Beaula J. T. Quantum chemical exploration on charge transfer interactions, vibrational, topological and molecular docking evaluation of the antimicrobial molecule 2,4, 6-triallyloxy-1,3,5-triazine. 2022. *Journal of Molecular Structure*, 1250 (2), P. 131846.
- [27]. Vang O., Ahmad N., Baile C.A., Baur J.A., Brown K., Csiszar A., Das D.K., Delmas D., Gottfried C., Lin H.Y., Ma Q.Y., Mukhopadhyay P., Nalini N., Pezzuto J.M., Richard T., Shukla Y., Surh Y.J., Szekeres T., Szkudelski T., Walle T., Wu J.M. What is new for an old molecule? Systematic review and recommendations on the use of resveratrol. 2011. *PLoS one*, 6(6), e19881.
- [28]. Jang, M., Cai, L., Udeani, G. O., Slowing, K. V., Thomas, C. F., Beecher, C. W. Cancer chemopreventive activity of resveratrol, a natural product derived from grapes. 1997. *Science*, 275, P. 218-220.
- [29]. Mazué F. Effects of red wine polyphenols on cell proliferation and on resveratrol metabolism. 2011. Doctoral thesis. Bourgogne University.
- [30]. Lançon, A., Frazzi, R. and Latruffe, N. Anti-oxidant, anti-inflammatory and anti-angiogenic properties of resveratrol in ocular diseases. 2016. *Molecules*, 21(3), P. 304.
- [31]. Schlienger J. L. Resveratrol and energy metabolism: a magic potion? *Medicine of Metabolic Diseases*, 2007. 1 (1): P. 77-78.
- [32]. Rong W., Weibiao Z. et Xiaohui J. «Reaction Kinetics of Degradation and Epimerization of Epigallocatechin Gallate (EGCG) in Aqueous System over a Wide Temperature Range». 2008. *Journal of Agricultural and Food Chemistry*, 56 (8): P. 2694-2701.
- [33]. Ahmed N.A., Radwan N.M., Aboul Ezz H.S., Salama N.A. The antioxidant effect of Green Tea Mega EGCG against electromagnetic radiation-induced oxidative stress in the hippocampus and striatum of rats. 2017. *Electromagnetic Biology and Medicine*, 36: P. 63–73.
- [34]. Rodriguez T. Pharmacological screening of the stem phenotype of ovarian cancer cells by epigallocatechin gallate. 2022. Master's Thesis in Biochemistry, Quebec University, P. 97.
- [35]. Yang Y., Zhao M., Liu Y., Fang Z., Li Q., Zheng, X. Yuerong L., Jianliang L., Separation and identification of an abundant trigalloylglucose from special tea genetic resources. 2022. *Beverage Plant Research*, 2 (1), P. 1-9.