

# Prospective observational study of drug prescribing patterns in elective surgeries: A hospital-based analysis

Ashif Ansari<sup>1</sup>, Sufiyan Khan<sup>2</sup>, Mohammad Saleem<sup>3</sup> Javed Ansari<sup>4\*</sup>, Pankaj kumar<sup>5</sup>

<sup>1</sup>. Department of Pharmacy Practice, Integral University, Lucknow, Uttar Pradesh, India.

<sup>2</sup>. Department of Pharmacy Practice, Integral University, Lucknow, Uttar Pradesh, India.

<sup>3</sup>. Faculty of Pharmacy, (Pharmacy Practice) Integral University, Lucknow, U.P, India.

<sup>4</sup>. Faculty of Pharmacy, Integral University, Lucknow, Uttar Pradesh, India.

<sup>5</sup>. Department of General Surgery, Integral Institute of Medical Science and Research Hospital, Lucknow, Uttar Pradesh, India.

\*Corresponding Author:- Javed Ansari

Collage full name-Faculty of Pharmacy, Integral University, Lucknow, Uttar Pradesh, India.

---

## Abstract

### Aim and Objectives:

Our study aims to evaluate the drug prescribing patterns in elective surgery in the Department of General Surgery at Integral Institute of Medical Sciences and Research, Integral University. This is a type of prospective observational study conducted using a sample size of 100 patients with different surgeries. The study span was a period of six months.

**Results:** A total of one hundred prescriptions were collected, and the medications prescribed for surgical patients were examined. Out of all 100 patients, men made up the majority of cases (59%) compared to women (41%). Ages 21 to 40 accounted for 52% of the cases. Cholelithiasis (30%) was the most common reason hospitalisation occurred, followed by hernias (19%). The most often prescribed medications were antimicrobials (21%), analgesics (11%), anti-ulcer (10%), and anti-emetic (4%) medication. Metronidazole (17%) and Cephalosporin (47%) were the two most often used antimicrobials. The most widely used analgesic was paracetamol (41.2%), which was followed by tramadol (34.2%). The most often prescribed medications from the anti-emetic and anti-ulcer categories were ondansetron (82%) and pantoprazole (77%).

**Conclusion:** This study, which included 100 patients, produced important information about the prescription practices for medications used in the surgery department. The kinds of information that people evaluate while deciding whether to have elective surgery vary widely. Patient case sheets are the most often used source of health information. Post-operative patients are frequently prescribed four or more medications, depending on the inclination and frequency. The majority of medications prescribed were GIT-related, AMAs, analgesics, multivitamins, and trace elements. Among antimicrobials, third-generation cephalosporins were the most frequently administered. The most commonly used analgesic was paracetamol. The anti-emetic and anti-ulcer medications that were most frequently administered were ondansetron and pantoprazole, respectively. Improvements in the prescribing pattern are required in accordance with worldwide norms; in particular, preventive prescribing of cephalosporin antibiotics and the use of generic medications and the Essential Medicines List (EML) to lower treatment costs are necessary. The most frequent reason for hospitalisation was cholelithiasis.

**Keywords:** Essential Medicine List, Post-Operative Medication, Antimicrobial Agents, Elective Surgery.

---

Date of Submission: 02-11-2024

Date of acceptance: 12-11-2024

---

## I. Introduction

Drug utilization study is “the marketing, distribution, prescription and use of a drug in society with special emphasis on resulting medical, social and economic consequences” [1]. Main aim of such study is to promote rational use of drugs in society. If a drug is prescribed in a rational way, it will help us in reducing the patient’s expenditure, lower adverse drug reactions, drug interactions and extra burden on medical as well as paramedical staff [2,3,4]. Therefore, prescription audit/monitoring or drug utilization study should be done

periodically to increase the therapeutic efficacy, decrease the adverse effects and provide feedback to the prescriber to ensure rational use of medicines. [5,6]

Surgery is a medical operation wherein a patient's tissues are cut or a previously incurred wound is closed. Surgery is the definition of treating bodily injuries or illnesses by incision or manipulation, often with tools [7]. Elective surgery, often called planned surgery is a term used to describe non-emergency surgery which is medically necessary, but which can be delayed for at least 24 hours [8]. Common examples of elective surgery include hip replacement, cataract extraction, ligament repairs, gall bladder surgery and hernia. There are three categories of elective surgery-

Non-urgent admission within 365 days is appropriate for a condition that is unlikely to deteriorate rapidly. Ex-Septoplasty.

Semi-urgent - Admission within 90 days is preferred for a condition that is unlikely to deteriorate fast. Ex-Colposcopy, amputation of the digit.

Urgent - Admission within 30 days is preferable for a condition with the potential to develop rapidly and become an emergency. Ex-Heart valve replacement, amputation of limb [9].

The study of drug prescribing trends in elective surgeries is an important undertaking that improves the quality and efficiency of healthcare services in the Department of General Surgery [10]. Present research focusses on which drugs are most regularly prescribed, why they are chosen, and how this information might assist patients and the healthcare system.

Such an analysis is essential to ensure patient safety and optimize healthcare outcomes. We can identify potential issues such as overuse, underuse, or inappropriate use of medications, thereby mitigating the risks associated with drug-related complications. Additionally, this research provides an evidence-based foundation for future clinical guidelines and protocols, ultimately benefiting both healthcare providers and patients [11].

## II. Methodology

The study was a six-month prospective observational study conducted in IPD and OPD departments General Surgery at Integral Institute of Medical Sciences and Research, Integral University. The CRF form was designed which includes the following information: name, age, gender, diagnosis, ongoing treatment of operation, drugs received by the patient, dosage, frequency of administration, and drugs recommended (generic/brand name) [12].

After interview of patient and review of case paper data regarding demographic details, clinical status, operative procedure, medication prescribed till discharge result of culture and sensitivity test, etc. were recorded in case record forms [13].

The study used a sample size of 100 patients who freely participated from general surgical wards and met the specified inclusion and exclusion criteria. Data collection during the study period was done manually [14]. Patients of either sex over 14 years of age, patients who had been through a surgery, patients with co-morbid condition, patients undergoing re-operation and patients visiting the OPD after surgery (day care surgery) were included in the study while patients who were below 14 years of age, pregnant and lactating women, cancer patients who were on chemotherapy and outpatients with minor ailments were excluded [15]. The data collected was analysed with the help of MS excel using descriptive statistics to determine drug use indicators and utilization pattern of drugs [16]. Ethical considerations were strictly followed, with complete adherence to WHO rules and permission from the Institutional Research and Ethics Committee (IEC/IIMSR/2023/46).

## III. Result

### Demographics (age and gender)

Total 100 cases were collected. Among 100 patients, 59 were males and 41 were females. This shows a higher proportion of male patients in comparison to female patients. Patients were grouped into 4 groups based on their age. Most of the cases noted were between 21-40 (52 cases) years of age followed by 41-60 (30 cases) years of age, 61-80 (12 cases) years of age. Least number of cases were found in <20 (6 cases) years of age. (Table 1).

Demographics	Frequency
<b>Gender</b>	
Male	59
Female	41
Total	100
<b>Age group (years)</b>	
<20	06
21-40	52
41-60	30

61-80	12
-------	----

**Table 1:** Demographics distribution of patients' gender and age.

### Disease conditions in general surgery

There were patients who had undergone surgical procedures of different diseases like cholelithiasis, hernia, fibroadenoma, appendicitis, hydrocele and haemorrhoid. Among these Cholelithiasis (30%) was the most common cause of hospitalisation followed by hernias (19%), fibroadenoma (13%), appendicitis (10%), haemorrhoid (8%) and hydrocele (6%). (**Table 2**).

Type of surgical procedure	Number of surgeries
Hernia	19
Cholelithiasis	30
Breast lumps/ Fibroadenoma	13
Appendicitis	10
Hydrocele	06
Hemorrhoid / Anal disease	08
Other	14
Total	<b>100</b>

**Table 2:** Disease conditions in general surgery

### Categories of drugs prescribed in general surgery

In 100 cases total 93 drugs of different classes have been prescribed and almost all prescription contains antimicrobial drugs (100%), Analgesics (96%), Anti-ulcer & anti-spasmodic (92%), multivitamin (88%), antiemetic (76%) and laxative (28%) followed by others (**Table 3-4**).

Parameters	Total no of prescription containing drug	Types of drugs
Antibiotics	100	20
Anti-ulcer and anti-spasmodic	92	09
Antiemetic	76	04
Laxative	28	03
Analgesics	96	10
Multivitamin	88	12
Others	70	35
TOTAL		93


**Table 3:** Drugs prescribed in general surgery patients.

Types of drugs prescribed	Total no of drugs	Percentage	Name of drugs
Anti-ulcer and anti-spasmodic	09	10	Pantoprazole, Rabeprazole, Ranitidine, Drotaverin, Hyoscine, Esomeprazole, Pepsin, Lansoprazole, Domperidone
Analgesics	10	11	Diclofenac, Aceclofenac, Tramadol, PCM, Ibuprofen, Naproxen, Chymotrypsin, celecoxib, Etoricoxib, Aspirin
Antibiotics	20	21	Ceftriaxone, Metronidazole, amoxicillin, Clavulanic acid, Amikacin, Clindamycin, Piperacillin + Tazobactam, Norfloxacin, Cefixime, Ciprofloxacin, Meropenem, Linezolid, Azithromycin, Doxycycline etc.
Multivitamin	12	13	Vitamin B complex, Vit. A, C, E, D, Zinc, Fatty acids, Carbohydrate
Antiemetics	03	04	Ondansetron, Metoclopramide, Domperidone
Laxatives	03	03	Lactulose, Liquid Paraffin, Milk of Magnesia
Others (Comorbidities)	35	38	Respiratory, DM, steroids, UTI, HTN, CAD related drugs

**Table 4:** Drugs Commonly Used in Surgical Patients

### Generic versus brand name prescription comparison


Out of 100 prescriptions 13 (13%) of the prescriptions are by their generic names and 87 (87%) by brand name. This shows a higher proportion of drug is prescribed by their brand name. **(Figure 1).**



**Figure 1:** Comparison of drugs prescribed by generic versus brand name.

### CATEGORIES OF ANTIMICROBIALS PRESCRIBED

The most commonly used antimicrobials drugs were Cephalosporin (47%), Penicillin (15%), Anti-amoebic (Metronidazole 17%) and others (21%). Among cephalosporin, first generation was used for about 2%, second generation 8.8%, third generation 83.6% and fourth generation about 5.4% **(Figure 2).**



**Figure 2:** Categories of antimicrobials drugs prescribed

### Combination of antibiotics drug prescribed


Fixed-dose combinations (FDCs) were given in which 1 Antibiotic was prescribed to 14 patients, 2 Antibiotics in 59, 3 Antibiotics in 23 and >3 Antibiotics in 04. Ceftriaxone was the most commonly used single antibacterial. Piperacillin + Tazobactam and Amoxicillin + Clavulanic acid was the most commonly used FDC combinations. This shows most of the prescriptions contain double antibiotics **(Table 5).**

Combination	Total no.
Single antibiotic	14
Double antibiotics	59
Triple antibiotics	23
More than three antibiotics	04

**Table 5:** Combination of antibiotics drug prescribed.

### Type of analgesics drug prescribed


In the study among analgesics, most commonly was used, Paracetamol (41.2%), Tramadol (34.2%), Diclofenac (10%), Chymotrypsin (7%), Aspirin (5.7%) and others (1.6%) **(Figure 3).**



**Figure 3:** Type of analgesics drug prescribed.

**Types of anti-emetics drug prescribed**


Most commonly used Anti-emetic were, Ondansetron (82%), Domperidone (13%) and Metoclopramide (5%) (Figure 4).



**Figure 4:** Types of anti-emetics drug prescribed

**Anti-ulcer drugs prescribed**

Anti-ulcer drugs prescribed mainly include, Pantoprazole (77%), Ranitidine (20%), Rabeprazole (2%) and Esomeprazole (1%) (Figure 5).



**Figure 5:** Anti-ulcer drugs prescribed.

#### **IV. Discussion**

Prescription reflects the physician's attitude towards the disease and treatment. It also provides a detailed image of the healthcare delivery system. Antibiotics are the primary treatment for infectious infections. In this study, we found that males accounted for 59% of the cases and females for 41%. Research found a similar preponderance [17-18]. The majority of the cases were between the ages of 21 and 40 (52 cases), followed by 41 to 60 (30 cases) and 61 to 80 (12 cases). The fewest cases were found among people aged 20 (6 cases).

In this study, disease patterns included cholelithiasis (30%), hernia (19%), breast lumps / fibroadenoma (13%), appendicitis (10%), haemorrhoids / anal disease (8%), and others (14%).

Antimicrobials (100%) were the most widely used medications in the study population, followed by analgesics (96%), anti-ulcerative (92%), anti-emetics (76%), multivitamins (88%), and others. The prescribing pattern for medications was consistent with other research [19].

Cephalosporins (47%) and metronidazole (17%) were the most widely used antimicrobials, followed by others. Other research reported similar findings. The most commonly utilised cephalosporins (83.6%) were third generation; comparable findings were reported in one investigation. For surgical prophylaxis, worldwide guidelines suggest using first- and second-generation cephalosporins rather than third- and fourth-generation cephalosporins, as well as other higher antimicrobials. The use of two antimicrobial drugs was also widespread, which corresponded to the findings reported by Khade et al. [20]. The use of one, two, and three antimicrobial agents was prevalent, in that order. The most often used antimicrobial combination was Ceftriaxone + Amikacin + Metronidazole, which was followed by Amoxicillin + Clavulanic acid, Piperacillin + Tazobactam, and Ceftriaxone + Amikacin + Metronidazole [20]. It was typical practice to combine penicillin with beta-lactamase inhibitors, then fluoroquinolones and third-generation cephalosporins. The older age group in this study is more involved, which increases the likelihood of morbidity. This supports the use of stronger, more general-purpose antibiotics as well.

Antibiotic overprescription and improper use raise the risk of drug resistance, adverse drug responses, and therapy costs. Parenteral route usage excessively is a prevalent practice in the majority of developing nations. This corresponds with the study's findings on all intravenous antibiotics. Few (13.4%) antibiotics were prescribed under their generic names, with the majority (86.6%) being prescribed under brand names [21]. Ceftriaxone was the most commonly used single antibacterial. Piperacillin + Tazobactam and Amoxicillin + Clavulanic acid was the most commonly used FDC combinations. Overall, beta lactam antibiotics, particularly Piperacillin were the most often administered antibiotics both pre- and post-surgical procedures [22]. Approximately 81% of the antimicrobials prescribed had approval from drug regulatory organisations and were listed in both the WHO-EML and NLEM [23].

#### **V. Limitations**

This study had a small sample size and was done in the patients admitted in the Department of Surgery. We did not enroll patients from Orthopedic, ENT, Eye/Ophthalmology, Obstetrics and Gynae Departments. Therefore, it gave us a limited pattern of drug use in post-operative patients in the surgical ward only. A prospective study, with large sample size in all the operative cases need to be done to evaluate the prescribing pattern and cost analysis in a better way.

#### **VI. Conclusion**

Most of the patients with surgical diseases require surgical intervention; therefore, they require more than one medication post-operatively to prevent wound infections and to relieve pain. There is a high tendency and frequency to prescribe more than four drugs, but less are prescribed from the National EML. Most of the drugs prescribed are generally AMAs, GIT related and Analgesics. There is an urgent need to develop proper prescription writing skills in budding doctors for the use of EML and generic medicines to reduce the cost of treatment and better compliance of the patients. The research carried out was an observational study conducted to assess the surgery diseases, surgical procedures and pattern of drug prescription. Patient with surgical diseases require more than one medication post operatively to prevent wound infection and to relieve pain. Cholelithiasis (30%) was the most common surgical disease noted, followed by Hernia (19%) and appendix (13%) was the most common surgical procedure done. Majority of drugs prescribed were mainly Antimicrobials (21%) followed by Analgesics (11%), Anti-ulcer (10%) and Antiemetic (9%). Among Antimicrobial agents, cephalosporin (47.5%) was mainly prescribed. Paracetamol (41.2%) was the most commonly used analgesic followed by Tramadol (34.2%), Ondansetron (82%) and Pantoprazole (77%) were the leading drugs used among anti emetic and anti-ulcer drugs. International guidelines indicate the use of 1st and 2nd generation cephalosporin rather than 3rd and 4th generation. There is a need to take steps to improve prescribing pattern as per international guidelines, especially cephalosporin antibiotics which is prescribed prophylactically.

## References

- [1]. World Health Organization. Introduction to Drug Utilization Research. Geneva: World health organization; 2003.
- [2]. Lfoholm PW, Katzung BG. Rational prescribing and prescription writing. In: Katzung BG, Masters SB, Trevor AJ, editors. Basic and Clinical Pharmacology. 11th Edition. New Delhi: Tata McGraw-Hill Education Private Limited.; 2009: 1127-36.
- [3]. World Health Organization. Teachers Guide to Good Prescribing. World Health Organization Department of Essential Drugs and Medicine Policy. Geneva: World Health Organization; 2001.
- [4]. World Health Organization. Introduction to Drug Utilization Research. Geneva: World Health Organization; 2003.
- [5]. Krishnaswamy K, Kumar BD, Radhaiah G. A drug survey – precepts and practices. *Eur J Clin Pharmacol.* 1985;29(3):363-70.
- [6]. Pradhan SC, Shewade DG, Shashindran CH, Bapna JS. Drug utilization studies. *Natl Med J India.* 1988;1:185-9.
- [7]. Kujath P, Michelsen A. Wounds–From physiology to wound dressing. *Deutsches Ärzteblatt International.* 2008 Mar;105(13):239.
- [8]. Curtis AJ, Wolfe R, Russell CO, et al. Determining priority for joint replacement: comparing the views of orthopedic surgeons and other professionals. *Med J Aust* 2011;195:699–702.
- [9]. SafiyaImtiazShaikh, D. Nagarekha, GanapatiHegade, and M. Marutheesh. Postoperative nausea and vomiting: A simple yet complex problem. *Anesth Essays Res.* 2016 Sep-Dec; 10(3): 388–396.
- [10]. Smith, J. (2019). *Pharmaceutical Management in Surgery*, 28(supplement 5), 168-172.
- [11]. Anderson MJ, DavidML, ScholzM, BullSJ, MorseD, Hulse-Stevens M, et al. Efficacy of skin and nasal povidone -iodine preparation against mupirocin resistant methicillin resistant staphylococcus aureus and s. aureus within the anterior antimicrobe agents chemother2015;59(5):2765e73
- [12]. Tefera BB, Getachew M, Kebede B. Evaluation of drug prescription pattern using World Health Organization prescribing indicators in public health facilities found in Ethiopia: systematic reviews and meta-analysis. *J Pharm Policy Pract.* 2021;14:1-10.
- [13]. World Health Organization. Introduction to Drug Utilization Research. 2003. Available at: <https://www.who.int/publications/i/item/8280820396>. Accessed on 15 May 2023.
- [14]. Ministry of Health and Family Welfare, Gov't of India. National List of Essential Medicines (NLEM). 2022. Available at: <https://main.mohfw.gov.in/?q=newshighlights-104>. Accessed on 15 May 2023.
- [15]. World Health Organization. WHO Model List of Essential Medicines – 23rd List. 2023. Available at: <https://www.who.int/publications/i/item/WHO-MHP HPS-EML-2023.02>. Accessed on 29 July 2023.
- [16]. Ministry of Health and Family Welfare Government of India. National List of Essential Medicines; 2015. Available at: <http://cdsco.nic.in/WriteReadData/NLEM 2015/NLEM, %202015.pdf>. Accessed 20 October 2016.
- [17]. Zeng G, Mai Z, Xia S, Wang Z, Zhang K, Wang L, et al. Prevalence of kidney stones in China: an ultrasonography based cross-sectional study. *BJU Int.* 2017;120(1):109-16.
- [18]. Faridi MS, Singh KS. Preliminary study of prevalence of urolithiasis in North-Eastern city of India. *J Fam Med Prim Care.* 2020;9(12):5939-43.
- [19]. Salman MT, Akram MF, Rahman S, Khan FA, Haseen MA, Khan SW. Drug Prescribing Pattern in Surgical Wards of a Teaching Hospital in North India. Vol. 5, No. 2 (2008-05 - 2008-06). ISSN : 0973 516X.
- [20]. A Khade, MSM Bashir, and A Sheethal. Prescription Pattern in the Department of Surgery in A Tribal District Hospital of Andhra Pradesh, India. *Ann Med Health Sci Res.* 2013 Jul Sep; 3(3): 438–441 .
- [21]. Liu M, Wu J, Xia W, Zhu Z, Chen H. Lifestyle factors, serum parameters, metabolic comorbidities, and the risk of kidney stones: a Mendelian randomization study. *Front Endocrinol.* 2023;14:1240171.
- [22]. Bratzler DW, Houck PM. Antimicrobial prophylaxis for surgery: An advisory statement from the National Surgical Infection Prevention Project. *Am J Surg.* 2005; 189:395–404.
- [23]. Kumar R et.al An in-depth study of drugs prescribing pattern in the Surgery Department of a Tertiary Care Teaching Institute in Northern India. *Int J Basic ClinPharmacol.*doi: 10.5455/2319 2003.ijbcp20140824, 2014 Aug;3(4):681-686